

Large-Population Passages of Vesicular Stomatitis Virus in Interferon-Treated Cells Select Variants of Only Limited Resistance

ISABEL S. NOVELLA,¹ MARTIN CILNIS,¹ SANTIAGO F. ELENA,^{2†} JOSHUA KOHN,¹ ANDRES MOYA,² ESTEBAN DOMINGO,³ AND JOHN J. HOLLAND^{1*}

Department of Biology and Institute for Molecular Genetics, University of California, San Diego, La Jolla, California 92093-0116,¹ and Departament de Genètica and Servei de Bioinformàtica, Facultat de Biologia, Universitat de València, 46100 Burjassot, València,² and Centro de Biología Molecular “Severo Ochoa” (Consejo Superior de Investigaciones Científicas, Universidad Autónoma de Madrid), Cantoblanco, 28049 Madrid,³ Spain

Received 27 March 1996/Accepted 6 June 1996

Vesicular stomatitis virus (VSV) populations were repeatedly passaged in L-929 cells treated with alpha interferon (IFN- α) at levels of 25 U/ml. This IFN- α concentration induced a 99.9% inhibition of viral yield in standard infections. Analysis of viral fitness (overall replicative ability measured in direct competition with a reference wild-type VSV) after 21 passages in IFN-treated cells showed only a limited increase or no increase in fitness, compared with the greater increase upon parallel passage in cells not treated with IFN- α . However, this limited increase in fitness was more pronounced when competition assays were carried out with IFN- α -treated cells, suggesting the selection of VSV populations with a low level of resistance to IFN- α . Thus, despite the extensively documented capacity of VSV to adapt to changing environments, the antiviral state induced by IFN- α imposes adaptive constraints on VSV which are not readily overcome.

Interferons (IFNs) are a group of cytokine proteins secreted by cells. They constitute one of the earliest responses after viral invasion, and they play a critical role in the clearance of some viral infections (31, 49). Gamma IFN (IFN- γ) and alpha and beta IFN (IFN- α and - β) act through different pathways to establish an antiviral state in cells stimulated by IFN binding to specific receptors. Many different antiviral proteins are induced by IFN, including the Mx proteins, the 2'-5'-oligoadenylate synthetase complex, and P1/eF-2 protein kinase (33, 52). IFN also upregulates the expression of major histocompatibility complex antigens (33, 52) and activates a double-stranded RNA adenosine deaminase (48). The editing function of the latter enzyme might hypermutate (and destabilize) RNA virus genomes (6). Thus, the antiviral actions of IFN are multiple, although the sensitivity of any particular virus to each of those mechanisms varies. Some DNA viruses carry accessory genes with anti-IFN activity (1, 2, 7, 12, 22, 34, 44, 58). RNA viruses have long been known for their potential to adapt to different environments (10, 11, 16, 26–29, 57). This adaptability is due mainly to their high mutation rates (16, 17, 29, 54), and their small genomic size does not usually allow the acquisition of genes mediating specific resistance to antiviral host functions. Despite this limitation, several RNA viruses have developed mechanisms for partial resistance to IFN. There are several reports of RNA virus proteins with anti-IFN activity (3, 24, 30, 32, 38, 50). Natural isolates of a particular virus often show variable sensitivity to IFN (4, 5, 21, 23, 36, 45, 46, 51), and patients undergoing IFN therapy often show different degrees of responsiveness to the treatment (23, 46). Understanding whether and how IFN influences RNA virus evolution is rele-

vant to its use in antiviral therapy, especially in the treatment of hepatitis B and C virus and human immunodeficiency virus infections (13, 25, 35, 37). In the present report, we describe the selection of vesicular stomatitis virus (VSV) populations displaying slight resistance to IFN. However, adaptation to growth in the presence of IFN also limits viral adaptation to the host cell type in which the viruses replicate repeatedly.

The Mudd-Summers (M-S) strain (Indiana serotype) of VSV was used in this study. The wild-type virus was a population derived from a clone, and it was partially adapted to mouse L-929 cells by four diluted passages before use. We have assigned a fitness value (W) of 1 (the neutral internal control value) to this wild-type VSV population. A relative fitness value is experimentally measured by coinfection passages of the test virus in direct competition with wild-type VSV. The test virus was a genetically marked (I1 monoclonal antibody resistant) subclone of wild-type VSV. The parental wild-type clone of VSV was employed as the competing reference virus. All virus assays were done in triplicate, both in the absence and in the presence of I1 monoclonal antibody in plaque overlay medium (27). The proportion of test and reference virus was measured initially (before passages) and after one or more competition passages. Changes in ratios of test to reference virus allowed the calculation of the relative fitness value (for further details, see references 8, 9, 18–20, and 27). It should be noted that in population biology, a fitness gain (or loss) of 10% is considered to be biologically highly meaningful (39). Two monoclonal antibody-resistant mutant (MARM) viral populations, MARM X and MARM E, were used in this study. We used a previously described clone, MARM X, which is an I1-antibody-resistant derivative of wild-type VSV. MARM X initially exhibited a low-fitness phenotype in L-929 cells ($W = 0.035 \pm 0.004$). The MARM E clone was obtained after 21 diluted passages of MARM X on L-929 monolayers and initially exhibited a high-fitness phenotype in that cell line ($W = 3.7 \pm 0.4$). Preparation of I1 monoclonal antibody stocks has been described elsewhere (20). Mouse IFN- α was obtained

* Corresponding author. Mailing address: Department of Biology and Institute for Molecular Genetics, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0116. Phone: (619) 534-2520. Fax: (619) 534-7108.

† Present address: Center for Microbial Ecology, Michigan State University, East Lansing, MI 48824-1325.

TABLE 1. Fitness changes of MARM X virus populations during large population passages in 25-U/ml IFN-treated and untreated L-929 cells^a

Series	Result with IFN in competition	Fitness (<i>W</i>)	<i>t</i> ^b	df	<i>P</i>
Control (untreated)					
A	-	3.1 ± 0.3			
	+	3.4 ± 0.1	0.6165	5	0.2823
B	-	3.4 ± 0.2			
	+	3.3 ± 0.3	0.2233	7	0.4148
C	-	0.89 ± 0.02			
	+	ND ^c			
D	-	2.6 ± 0.3			
	+	2.2 ± 0.1	1.1461	7	0.1447
E	-	4.6 ± 0.8			
	+	4.3 ± 0.7	0.3255	7	0.3772
F	-	2.5 ± 0.1			
	+	2.44 ± 0.08	0.5005	7	0.3160
IFN-treated					
B	-	0.31 ± 0.07			
	+	1.02 ± 0.03	9.0879	7	<0.0001
C	-	0.47 ± 0.08			
	+	1.1 ± 0.6	0.9762	8	0.1788
D	-	0.5 ± 0.1			
	+	0.9 ± 0.3	1.3700	8	0.1040
E	-	0.49 ± 0.09			
	+	5 ± 1	3.7908	7	0.0034
F	-	1.0 ± 0.2			
	+	2.9	2.1347	8	0.0327

^a The results were analyzed by ANCOVA (55). Parental MARM X has a fitness of 0.035 ± 0.004 in untreated L-929 cells, and fitness values in the presence of IFN are not statistically different ($P = 0.414$).

^b Student's *t* test employed for comparison of each population's competition in the presence and absence of IFN.

^c ND, not determined.

from Lee Biomolecular Research, Inc. Twenty-one consecutive large virus population (control) transfers in the absence of IFN were done as described previously (8, 9, 18–20, 27). Also, 21 passages in IFN-pretreated cells were carried out in the same manner. Virus yields were diluted after each passage so that approximately 10^5 to 10^6 infectious particles were present in each transferred population. Cells were pretreated with 1.5 ml of a 25-U/ml IFN solution for 24 h. After pretreatment, IFN was removed and the monolayers were immediately infected with the large virus populations. IFN inhibition of virus yield was about 1,000-fold (99.9% yield reduction). Viral titers were determined after each passage, so that appropriate dilutions were used in the next passage to keep the population transfer size approximately constant (10^5 to 10^6 infectious particles) at a multiplicity of infection of about 0.1.

MARM X populations passaged 21 times in IFN-treated cells showed significant increases in mean fitness in untreated L-929 cells ($W = 0.51 \pm 0.06$; $P < 0.0001$) compared with the level in the control parental population ($W = 0.035 \pm 0.04$), as shown by an analysis of covariance (ANCOVA) test (55). However, such increases were significantly lower than those observed in populations passaged in the absence of IFN ($W = 2.3 \pm 0.3$; $P < 0.0001$). Fitness gains of populations passaged in IFN-treated cells were higher when competitions were carried out in the IFN-treated cells than in the untreated cells (1.7 ± 0.4 versus 0.51 ± 0.06 ; $P = 0.001$) (Table 1). This difference indicated the selection of populations with slightly increased resistance to IFN. Results with MARM E populations were less significant (Table 2). First, MARM E populations passaged in IFN-treated cells did not show significant

overall increases in fitness in L-929 cells ($W = 6.0 \pm 1$) compared with unpassaged MARM E populations ($W = 3.7 \pm 0.3$). The results of each series were very heterogeneous, with values higher than, lower than, or about equal to those for parental MARM E (Table 2). The possible generation of an IFN-resistant population was statistically analyzed by ANCOVA ($P = 0.0525$) and ANOVA ($P = 0.0233$). Both results are close to the 0.05 level of significance, indicating that only very limited resistance was developed in MARM E populations.

Next, we carried out experiments with cells pretreated with low concentrations of IFN (1.5 ml of a 1-U/ml solution for 6 h, causing about 60% inhibition of viral yield). Neither MARM X nor MARM E virus showed any evidence of resistance after 21 passages at low levels of IFN, and their adaptation to L-929 cells was comparable to that observed in untreated controls in every case (data not shown).

The present study shows that, at best, only partially IFN-resistant variants of VSV can be selected even when a strong selective pressure is applied. This very limited acquisition of IFN- α resistance is probably due to the pleiotropic effects of IFN: inhibition of protein synthesis, RNA degradation by endonucleases, RNA editing, and inhibition of viral entrance in the case of VSV (56, 59). The multiple effects of IFN apparently severely constrain the possibility of generating highly resistant VSV genomes. Presumably, multiple mutations at several genomic loci would be needed to overcome the multiple biochemical blocks in IFN-treated cells. This would be a low-probability occurrence in our viral quasispecies populations (15). VSV is known to be very sensitive to IFN, and it is

TABLE 2. Changes in fitness of MARM E virus populations during large population passages in 25-U/ml IFN-treated and untreated L-929 cells^a

Series	Result with IFN in competition	Fitness (<i>W</i>)	<i>t</i> ^b	df	<i>P</i>
Control (untreated)					
A	-	9 ± 2			
	+	11 ± 3	0.3996	8	0.35
B	-	13 ± 2			
	+	11.2 ± 0.3	0.5646	8	0.2939
C	-	19 ± 3			
	+	8.7 ± 0.8	3.9085	8	0.0022
D	-	11 ± 2			
	+	15 ± 3	0.9631	8	0.1819
E	-	16 ± 3			
	+	15 ± 2	0.1488	6	0.4433
F	-	28 ± 2			
	+	21 ± 3	1.7442	8	0.0596
IFN treated					
A	-	17 ± 2			
	+	18 ± 8	0.0811	8	0.4687
B	-	8.5 ± 0.2			
	+	31 ± 15	1.527	8	0.0826
C	-	24 ± 8			
	+	28 ± 15	0.2068	7	0.4210
D	-	4 ± 1			
	+	21 ± 5	3.3826	8	0.0048
E	-	3.0 ± 0.9			
	+	14 ± 5	2.2025	7	0.0317
F	-	0.7 ± 0.2			
	+	3.7 ± 0.4	6.1426	6	0.0004

^a Results were analyzed by ANCOVA (55). Parental MARM E has a fitness of 3.7 ± 0.3 in untreated L-929 cells, and fitness values in the presence of IFN are not statistically different ($P = 0.483$).

^b Student's *t* test employed for comparison of each population's competition in the presence and absence of IFN.

routinely used to test IFN activity, indicating a lack of genetic functions conferring IFN resistance. In some cases, as described for mengo virus, resistant phenotypes are simply explained by the inability of a viral strain to induce IFN in infected cells (40). VSV Indiana can induce IFN production in vivo (60), but not in cell culture (43). In fact, VSV Indiana strains are often suppressors of IFN production in cell culture (41–43, 53). However, in our case, IFN is supplied exogenously; thus, our resistant populations are not due to variants that have lost the capacity to induce IFN.

In summary, despite their great adaptability and remarkable ability to gain fitness exponentially (47), RNA viruses have only a feeble capacity to increase fitness in the presence of IFN. Furthermore, this slight gain in fitness in the presence of IFN is achieved only at the cost of an overall decrease in gains of fitness in the host cell. Our results demonstrate that the adaptability of RNA viruses encounters insurmountable limitations when antiviral activities are directed to multiple viral functions and/or to independent targets on the viral genome. They provide further support for the notion that whole-virus vaccines able to trigger a multivalent immune response or combination therapy with multiple antiviral inhibitors represents the most logical approach for the control of highly variable RNA viruses (14, 15).

We thank Estelle Bussey for excellent technical assistance, Leo Lefrancoise for kindly providing I1 hybridoma cells, and Philip Marcus for helpful discussions.

This research was supported by grant DGICYT PB94-0034 (Madrid and Valencia), grant FIS 95/0034-01 and Fundacion Ramon Areces (Madrid), and NIH grant AI14627. S.F.E. was supported by a fellowship from Conselleria d'Educacio y Ciencia (Generalitat Valenciana).

REFERENCES

- Akkrāju, G. R., P. Whitaker-Dowling, J. S. Youngner, and R. Jagus. 1989. Vaccinia specific kinase inhibitory factor prevents translational inhibition by double-stranded RNA in rabbit reticulocyte lysate. *J. Biol. Chem.* **264**: 10321–10325.
- Alcami, A., and G. L. Smith. 1995. Cytokine receptors encoded by poxviruses: a lesson in cytokine biology. *Immunol. Today* **16**:474–478.
- Beattie, E., K. L. Denzler, J. Tartaglia, M. E. Perkus, E. Paoletti, and B. L. Jacobs. 1995. Reversal of the interferon-sensitive phenotype of a vaccinia virus lacking E3L by expression of the reovirus S4 gene. *J. Virol.* **69**:499–505.
- Carrigan, D. R., and C. M. Kabacoff. 1987. Identification of a nonproductive cell-associated form of measles virus by its resistance to inhibition by recombinant human interferon. *J. Virol.* **61**:1919–1926.
- Carrigan, D. R., and K. K. Knox. 1990. Identification of interferon-resistant subpopulations in several strains of measles virus: positive selection by growth of the virus in brain tissue. *J. Virol.* **64**:1606–1615.
- Cattaneo, R., and M. Billeter. 1992. Mutations and A/I hypermutations in measles virus persistent infections. *Curr. Top. Microbiol. Immunol.* **176**:63–74.
- Cayley, P. L., J. A. Davies, K. G. McCullagh, and I. M. Kerr. 1984. Activation of the pp(A2')nA system in interferon-treated herpes simplex virus-infected cells and evidence for novel inhibitors of the ppp(A2')n-AA-dependent RNase. *Eur. J. Biochem.* **143**:165–174.
- Clarke, D. K., E. A. Duarte, S. Elena, A. Moya, E. Domingo, and J. Holland. 1994. The red queen reigns in the kingdom of RNA viruses. *Proc. Natl. Acad. Sci. USA* **91**:4821–4824.
- Clarke, D. K., E. A. Duarte, A. Moya, S. F. Elena, E. Domingo, and J. Holland. 1993. Genetic bottlenecks and population passages cause profound fitness differences in RNA viruses. *J. Virol.* **67**:222–228.
- Coffin, J. M. 1990. Genetic variation in retroviruses. *Appl. Virol. Res.* **2**:11–33.
- Coffin, J. M. 1992. Genetic diversity and evolution of retrovirus. *Curr. Top. Microbiol. Immunol.* **176**:143–164.
- Davies, M. V., O. Elroy-Stein, R. Jagus, B. Moss, and R. J. Kaufman. 1992. The vaccinia virus K3L gene product potentiates translation by inhibiting double-stranded-RNA-activated protein kinase and phosphorylation of the alpha subunit of eukaryotic initiation factor 2. *J. Virol.* **66**:1943–1950.
- DiBisceglie, A. M., P. Martin, C. Kassianides, M. Lisker-Melamm, L. Murray, J. Waggoner, Z. Goodman, S. M. Banks, and J. H. Hoofnagle. 1989. Recombinant interferon alfa therapy for chronic hepatitis C. A randomized, double-blind, placebo-controlled trial. *N. Engl. J. Med.* **321**:1506–1510.
- Domingo, E. 1989. RNA virus evolution and the control of viral disease. *Prog. Drug Res.* **33**:93–133.
- Domingo, E., and J. J. Holland. 1992. Complications of RNA heterogeneity for the engineering of virus vaccines and antiviral agents, p. 13–31. *In* J. K. Setlow (ed.), *Genetic engineering, principles and methods*. Plenum Press, New York.
- Domingo, E., and J. J. Holland. 1994. Mutation rates and rapid evolution of RNA viruses, p. 161–183. *In* S. Morse (ed.), *The evolutionary biology of viruses*. Raven Press, New York.
- Drake, J. W. 1993. Rates of spontaneous mutations among RNA viruses. *Proc. Natl. Acad. Sci. USA* **90**:4171–4175.
- Duarte, E., D. Clarke, A. Moya, E. Domingo, and J. Holland. 1992. Rapid fitness losses in mammalian RNA virus clones due to Muller's ratchet. *Proc. Natl. Acad. Sci. USA* **89**:6015–6019.
- Duarte, E. A., D. C. Clarke, A. Moya, S. F. Elena, E. Domingo, and J. Holland. 1993. Many-trillionfold amplification of single RNA virus particles fails to overcome the Müller's ratchet effect. *J. Virol.* **67**:3620–3623.
- Duarte, E. A., I. S. Novella, S. Ledesma, D. C. Clarke, A. Moya, S. F. Elena, E. Domingo, and J. J. Holland. 1994. The subclonal components of consensus fitness in an RNA virus clone. *J. Virol.* **68**:4295–4301.
- Enomoto, N., M. Kurosaki, Y. Tanaka, F. Marumo, and C. Sato. 1994. Fluctuation of hepatitis C virus quasiespecies in persistent infection and interferon treatment revealed by single-strand conformation polymorphism analysis. *J. Gen. Virol.* **75**:1361–1369.
- Esteban, M., J. Benavente, and E. Paez. 1986. Mode of sensitivity and resistance of vaccinia virus replication to interferon. *J. Gen. Virol.* **67**:801–808.
- Gil, B., C. Qian, J. I. Riezu-Boj, M. P. Civeira, and J. Prieto. 1993. Hepatic and extrahepatic HCV RNA strands in chronic hepatitis C: different patterns of response to interferon treatment. *Hepatology* **18**:1050–1054.
- Gunnery, S., A. P. Rice, H. D. Robertson, and M. B. Mathews. 1990. Tat-responsive region RNA of human immunodeficiency virus 1 can prevent activation of the double-stranded-RNA-activated protein kinase. *Proc. Natl. Acad. Sci. USA* **87**:8687–8691.
- Gutterman, J. U. 1994. Cytokine therapeutics: lessons from interferon α . *Proc. Natl. Acad. Sci. USA* **91**:1198–1205.
- Holland, J. J. 1992. Genetic diversity of RNA viruses. Springer-Verlag, Berlin.
- Holland, J. J., J. C. de la Torre, D. K. Clarke, and E. Duarte. 1991. Quantitation of relative fitness and great adaptability of clonal populations of RNA viruses. *J. Virol.* **65**:2960–2967.
- Holland, J. J., J. C. de la Torre, and D. Steinhauer. 1992. RNA virus populations as quasiespecies. *Curr. Top. Microbiol. Immunol.* **176**:1–20.
- Holland, J. J., K. Spindler, F. Horodyski, E. Grabau, S. Nichol, and S. VandePol. 1982. Rapid evolution of RNA genomes. *Science* **215**:1577–1585.
- Imani, F., and B. L. Jacobs. 1988. Inhibitory activity for the interferon-induced protein kinase is associated with the reovirus serotype 1 $\sigma 3$ protein. *Proc. Natl. Acad. Sci. USA* **85**:7887–7891.
- Karupiah, G., T. N. Fredrickson, K. L. Holmes, L. H. Khairallah, and R. M. L. Buller. 1993. Importance of interferons in recovery from mousepox. *J. Virol.* **67**:4214–4226.
- Katze, M. G., B. M. Detjen, B. Safer, and R. M. Krug. 1986. Translational control by influenza virus: suppression of the kinase that phosphorylates the alpha subunit of initiation factor eIF-2 and selective translation of influenza viral mRNAs. *Mol. Cell. Biol.* **6**:1741–1750.
- Kerr, I. M., and G. R. Stark. 1992. The antiviral effects of interferon and their inhibition. *J. Interferon Res.* **12**:237–240.
- Kitajewski, J., R. J. Schneider, B. Safer, S. M. Munemitsu, C. E. Samuel, B. Thimmappaya, and T. Shenk. 1986. Adenovirus VAI RNA antagonizes the antiviral action of interferon by preventing activation of the interferon-induced eIF-2 α protein kinase. *Cell* **45**:195–200.
- Koziel, M. J., and B. D. Walker. 1992. Viruses, chemotherapy and immunity. *Parasitology* **105**:s85–s92.
- Kunzi, M. S., H. Farzadegan, J. B. Margolock, D. Vlahov, and P. M. Pitha. 1995. Identification of human immunodeficiency virus primary isolates resistant to interferon α and correlation to prevalence to disease progression. *J. Infect. Dis.* **171**:822–828.
- Lane, H. C., J. Feinberg, V. Davey, L. Deyton, M. Baseler, J. Manischewitz, H. Masur, J. A. Kovacs, B. Herpin, R. Walker, J. A. Metcalf, N. Salzman, G. Quinann, and A. S. Fauci. 1988. Anti-retroviral effects of interferon- α in AIDS-associated Kaposi's sarcoma. *Lancet* **ii**:1218–1222.
- Lee, T. G., J. Tomita, A. G. Hovanessian, and M. G. Katze. 1990. Purification and partial characterization of a cellular inhibitor of interferon-induced protein kinase of Mr 68,000 from influenza virus-infected cells. *Proc. Natl. Acad. Sci. USA* **87**:6208–6212.
- Lewontin, R. C. 1974. *The genetic basis of evolutionary change*. Columbia University Press, New York.
- Marcus, P. I., P. T. Guidon, Jr., and M. J. Sekellick. 1981. Interferon induction by viruses. VII. Mengovirus: "interferon sensitive" mutant phenotype attributed to interferon-inducing particle activity. *J. Interferon Res.* **1**:601–611.
- Marcus, P. I., and M. J. Sekellick. 1985. Interferon induction by viruses. XII.

- Detection and assay of interferon induction-suppressing particles of vesicular stomatitis virus. *Virology* **142**:411–415.
42. **Marcus, P. I., and M. J. Sekellick.** 1987. Interferon induction by viruses. XV. Biological characteristics of interferon induction-suppressing particles of vesicular stomatitis virus. *J. Interferon Res.* **7**:269–284.
 43. **Marcus, P. I., M. J. Sekellick, and S. T. Nichol.** 1992. Interferon induction by viruses. XXI. Vesicular stomatitis virus: interferon inducibility as a phylogenetic marker. *J. Interferon Res.* **12**:297–305.
 44. **Mellits, K. H., M. Kostura, and M. B. Mathews.** 1990. Interaction of adenovirus VA RNA1 with the protein kinase DAI: nonequivalence of binding and function. *Cell* **61**:843–852.
 45. **Moskophidis, D., M. Bategay, M. A. Bruendler, E. Laine, I. Gresser, and R. M. Zinkernagel.** 1994. Resistance of lymphocytic choriomeningitis virus to alpha/beta interferon and to gamma interferon. *J. Virol.* **68**:1951–1955.
 46. **Naoumov, N. V., M. G. Thomas, A. L. Mason, S. Chokshi, C. J. Bodicky, F. Farzaneh, R. Williams, and R. P. Perrillo.** 1995. Genomic variations in the hepatitis B core gene: a possible factor influencing response to interferon alfa treatment. *Gastroenterology* **108**:505–514.
 47. **Novella, I. S., E. A. Duarte, S. F. Elena, A. Moya, E. Domingo, and J. J. Holland.** 1995. Exponential increases of RNA virus fitness during large population transmissions. *Proc. Natl. Acad. Sci. USA* **92**:5841–5844.
 48. **Patterson, J. B., and C. E. Samuel.** 1995. Expression and regulation by interferon of a double-stranded RNA-specific adenosine deaminase from human cells: evidence for two forms of the deaminase. *Mol. Cell. Biol.* **15**:5376–5388.
 49. **Riviere, Y., I. Gresser, J. C. Guillon, and M. G. Tovey.** 1977. Inhibition by anti-interferon serum of lymphocytic choriomeningitis virus disease in suckling mice. *Proc. Natl. Acad. Sci. USA* **74**:2135–2139.
 50. **Roy, S., M. G. Katze, N. T. Parkin, I. Ederly, A. G. Hovanessian, and N. Sonenberg.** 1990. Control of the interferon-induced 68-kilodalton protein kinase by the HIV-1 *tat* gene product. *Science* **247**:1216–1219.
 51. **Saitoh, H., S. Naitoh, H. Okamoto, and Y. Akahane.** 1994. Prompt decrease of circulating hepatitis C virus in patients with chronic hepatitis C after treatment with interferon. *J. Interferon Res.* **14**:239–244.
 52. **Samuel, C. E.** 1991. Antiviral actions of interferon. Interferon-regulated cellular proteins and their surprising selective antiviral activities. *Virology* **183**:1–11.
 53. **Sekellick, M. J., and P. I. Marcus.** 1989. Interferon induction by viruses. XVII. Non-temperature-sensitive mutations regulate interferon induction by vesicular stomatitis virus. *J. Gen. Virol.* **70**:405–415.
 54. **Smith, D. B., and S. C. Inglis.** 1987. The mutation rate and variability of eukaryotic viruses: an analytical review. *J. Gen. Virol.* **68**:2729–2740.
 55. **Sokal, R. R., and F. J. Rohlf.** 1981. *Biometry*. Freeman, New York.
 56. **Vilcek, J., and G. C. Sen.** 1996. Interferons and other cytokines, p. 375–399. *In* B. N. Fields, D. M. Knipe, and P. M. Howley (ed.), *Virology*. Lippincott-Raven Publishers, Philadelphia.
 57. **Wain-Hobson, S.** 1992. Human immunodeficiency virus type 1 quasispecies *in vivo* and *ex vivo*. *Curr. Top. Microbiol. Immunol.* **176**:181–194.
 58. **Whitaker-Dowling, P., and J. S. Youngner.** 1984. Characterization of a specific kinase factor produced by vaccinia virus which inhibits the interferon-induced protein kinase. *Virology* **137**:171–181.
 59. **Whitaker-Dowling, P. A., D. K. Wilcox, C. C. Widnell, and J. S. Youngner.** 1983. Interferon-mediated inhibition of virus penetration. *Proc. Natl. Acad. Sci. USA* **80**:1083–1086.
 60. **Youngner, J. S., and G. Wertz.** 1968. Interferon induction in mice by vesicular stomatitis virus. *J. Virol.* **2**:1360–1361.