

ERRATUM

Characterization of Entry Mechanisms of Human Herpesvirus 8 by Using an Rta-Dependent Reporter Cell Line

Naoki Inoue, Jörn Winter, Renu B. Lal, Margaret K. Offermann, and Shin Koyano

*Division of Viral and Rickettsial Diseases and Division of AIDS, STD, TB and Laboratory Research, Centers
for Disease Control and Prevention, and Winship Cancer Center, Emory University, Atlanta, Georgia*

Volume 77, no. 14, p. 8147–8152, 2003. Page 8149, legend to Fig. 2, lines 8 to 12: “The cells were also incubated in the presence of the indicated amount of FN-like polymer and RGD peptide at 4°C (B) or in the presence of soluble $\alpha 3\beta 1$ and $\alpha 5\beta 1$ integrins at 37°C (C) for 90 min, washed with the inhibitor-free medium, and then incubated with the HHV-8 inoculums for 2 h (B, C)” should read “The cells were also incubated in the presence of the indicated amount of FN-like polymer and RGD peptide at 4°C for 90 min, washed with the inhibitor-free medium, and then incubated with the HHV-8 inoculums for 2 h (B). The cells were incubated for 2 h with HHV-8 inoculums that had been preincubated in the presence of $\alpha 3\beta 1$ and $\alpha 5\beta 1$ integrins at 37°C for 90 min (C).”