

Complete DNA Sequences of Two Oka Strain Varicella-Zoster Virus Genomes[∇]

Sueli L. Tillieux,¹ Wendy S. Halsey,² Elizabeth S. Thomas,² John J. Voycik,^{2,‡}
Ganesh M. Sathe,² and Ventsislav Vassilev^{1,*}

GlaxoSmithKline Biologicals, Research and Development, Viral Vaccines, B-1330 Rixensart, Belgium,¹ and GlaxoSmithKline, Discovery Technology Group, 1250 South Collegeville Road, Collegeville, Pennsylvania 19426²

Received 10 April 2008/Accepted 3 September 2008

Varicella-zoster virus (VZV) is a herpesvirus and is the causative agent of chicken pox (varicella) and shingles (herpes zoster). Active immunization against varicella became possible with the development of live attenuated varicella vaccine. The Oka vaccine strain was isolated in Japan from a child who had typical varicella, and it was then attenuated by serial passages in cell culture. Several manufacturers have obtained this attenuated Oka strain and, following additional passages, have developed their own vaccine strains. Notably, the vaccines Varilrix and Varivax are produced by GlaxoSmithKline Biologicals and Merck & Co., Inc., respectively. Both vaccines have been well studied in terms of safety and immunogenicity. In this study, we report the complete nucleotide sequence of the Varilrix (Oka-V_{GSK}) and Varivax (Oka-V_{Merck}) vaccine strain genomes. Their genomes are composed of 124,821 and 124,815 bp, respectively. Full genome annotations covering the features of Oka-derived vaccine genomes have been established for the first time. Sequence analysis indicates 36 nucleotide differences between the two vaccine strains throughout the entire genome, among which only 14 are involved in unique amino acid substitutions. These results demonstrate that, although Oka-V_{GSK} and Oka-V_{Merck} vaccine strains are not identical, they are very similar, which supports the clinical data showing that both vaccines are well tolerated and elicit strong immune responses against varicella.

Varicella-zoster virus (VZV) is a human alphaherpesvirus that causes chicken pox (varicella) and shingles (herpes zoster) (75). VZV has a linear, double-stranded DNA genome of approximately 125 kb that encodes at least 71 proteins (12). Primary infection with VZV results in varicella, which is a widespread, highly contagious disease. Varicella is commonly regarded as a mild childhood illness, but it may lead to serious complications, such as secondary bacterial infection, pneumonia, encephalitis, congenital infection, and death (76).

Like other herpesviruses, VZV has the capacity to persist in the body after the primary acute infection as a latent infection in sensory nerve ganglia. This lifelong latent infection commonly reactivates to cause herpes zoster, typically in elderly or immunocompromised patients (65).

In 1974, Takahashi et al. reported the development of a live-attenuated varicella vaccine through serial passages of wild-type virus in cell culture (67). The parental virus, Oka-P, was isolated in primary human embryo lung cell culture from vesicle fluid from a 3-year-old boy with typical varicella. The virus was attenuated by 10 passages in HEL and 12 passages in guinea pig embryo cells, plaque-purified, and passaged five times further in human diploid cells (WI38) to prepare a strain suitable for use as a vaccine (Oka-V) (Fig. 1) (67).

The Oka-V strain was first supplied in 1976 under license

from the Biken Institute in Japan. Several manufacturers (SmithKline RIT, Merck Sharp & Dohme, and Pasteur Mérieux) subsequently used the Oka-V strain in the development of proprietary vaccines. A product license was obtained for Varilrix (frozen formulation) in 1984 by SmithKline RIT for use in groups at high risk for severe varicella and their healthy close contacts. SmithKline RIT—now GlaxoSmithKline (GSK) Biologicals—subsequently developed a refrigerator-stable formulation of this varicella vaccine. Varilrix is indicated in many countries for use in healthy and immunocompromised subjects from 9 months of age. GSK Biologicals' varicella vaccine production is based on the seed lot system (6, 14) using classical cell culture methods (Fig. 1). A manufacturer's working cell bank of human diploid cells, MRC-5, was prepared and tested according to World Health Organization requirements.

The Biken vaccine was licensed in Japan and Korea, in 1986 and 1988, respectively, for use in healthy subjects, and a license for Varivax with the same indication was granted in the United States in 1995 (1). In 1993, the vaccine manufactured by Pasteur Mérieux was licensed in France for use in potentially immunocompromised subjects.

Although the varicella vaccine is licensed in many countries, it is not routinely used because complications associated with varicella disease are often underestimated. Universal mass vaccination against varicella is implemented only in few countries; however, it is under consideration in many others (38, 40, 54, 72). The incidence of varicella disease and the rate of varicella-related hospitalizations in the United States have declined by about 80% since implementation of universal mass vaccination against varicella (using Varivax) in 1996 (8, 16, 81). A similar

* Corresponding author. Mailing address: Viral Vaccines Research and Development, GlaxoSmithKline Biologicals, Rue de l'Institut, 89 (P31-005), B-1330 Rixensart, Belgium. Phone: 32-2-656-6422. Fax: 32-2-656-8113. E-mail: ventsislav.vassilev@gskbio.com.

‡ Present address: PerkinElmer Life and Analytical Sciences, 710 Bridgeport Ave, Shelton, CT 06484-4794.

[∇] Published ahead of print on 10 September 2008.

OSAKA UNIVERSITY	Clinical case of varicella (1972)	Virus passage	(Oka-P)
	Isolation on HEL	1	
	10 passages on HEL	11	
	12 passages on GPE	23	
	5 passages on WI38		
	Master Seed Lot	28	(Oka-V)
GSK BIOLOGICALS	4 passages on MRC-5		
	GSK Biologicals		
	Working Seed Lots	32	
	3 passages on MRC-5		
Varicella inoculum	35		
	3 passages on MRC-5		
	Varicella bulk vaccine	38	(Oka-V _{GSK})

FIG. 1. Passage history of the live attenuated Oka varicella vaccine. HEL, human embryonic lung cells; GPE, guinea pig embryo cells; WI38 and MRC-5, human diploid cells.

decrease was observed in Uruguay since the introduction of varicella vaccination (using Varilrix) into the routine childhood immunization program in 1999, with the greatest reduction in children aged 1 to 4 years (51). Most pre- and postlicense studies showed that vaccination with one dose of varicella-containing vaccine provides 70% to 90% protection from chicken pox and over 95% protection against the most severe forms of the disease for a 7- to 10-year period after vaccination (2, 17, 33, 33a, 34, 40, 61). However, vaccine-induced immunity wanes over time (9), leading countries such as the United States to recommend a two-dose schedule for varicella vaccination (40). This strategy aims to overcome primary vaccine failures and to improve long-term protection, thereby reducing the risk of breakthrough varicella (4). Combined vaccine products containing the VZV Oka strain have been developed as well. For instance, GSK Biologicals and Merck & Co., Inc., developed combined tetravalent measles-mumps-rubella-varicella vaccines (Priorix-Tetra and ProQuad, respectively), providing the benefits of measles-mumps-rubella and varicella vaccination in a single injection (19, 30, 35, 48, 71, 72, 79).

Different sets of serological readouts have been used to characterize the adaptive humoral immune response after varicella vaccination or infection (4, 13, 26, 31, 34, 58, 73, 74). Comparative analysis has raised the possibility that differences in the genetic code between the vaccine strains could be responsible for disparity in vaccine-induced humoral responses (36).

Oka-V, and presumably its derivative vaccine strains, was not cloned during the development and the preparation of vaccine (67). Sequencing of the complete genome of the orig-

inal Oka-V vaccine preparation revealed that it contained multiple variants that could be separated in cell culture (20, 22).

The aim of the present study was to analyze the complete consensus nucleotide sequences of Oka-V strain viruses contained in Varilrix (GSK Biologicals; Oka-V_{GSK}) and Varivax (Merck & Co., Inc.; Oka-V_{Merck}) and to compare them to the published sequences of Oka-V and Oka-P (22). The full-length genomic sequences were also compared to published partial sequencing information on Oka-V_{GSK} and Oka-V_{Merck} (3, 32, 60, 63).

MATERIALS AND METHODS

Nucleic acid extraction. Total DNA was extracted from a single vial of recent production lots of Varilrix (lot VAV10118, produced in April 2002; GSK Biologicals, Rixensart, Belgium) and Varivax (lot 0895 M, purchased in 2003; Merck & Co., Inc., Whitehouse Station, NJ) vaccines using a High Pure viral nucleic acid kit from Roche (Basel, Switzerland). In brief, 100 μ l of sample was lysed in a lysing-binding buffer in the presence of proteinase K. The lysis mixture was then applied to a glass fiber filter, which binds the nucleic acids in the presence of the lysis and binding buffer containing chaotropic salts. Bound nucleic acids were eluted in 50 μ l of nuclease-free water by centrifugation and stored at -70°C .

PCR. Around 540 primers were designed using Primer D software (GSK in-house software) and the nucleotide sequence of the Dumas strain (GenBank accession no. X04370) (12). Overlapping primers were designed approximately 500 bases apart to cover the entire genomic sequence of VZV. Sequences of primers used for amplification and sequencing are available upon request. The reaction mixtures for PCR contained 15 μ l of HotStarTaq Plus Master Mix solution (Qiagen, Valencia, CA), 0.3 μ M of each primer, and 5 ng of template DNA. A Tetrad thermal cycler (MJ Research, Waltham, MA) was used for all amplifications. An initial hot-start PCR step of 96°C for 15 min was followed by 35 cycles of amplification (95°C for 20 s, 55°C for 30 s, and 72°C for 45 s) and a final elongation step at 72°C for 3 min. All amplified products were then purified using QIAquick PCR purification kit (Qiagen). Direct sequencing of both DNA strands was performed on the generated amplicons.

Sequencing. Direct sequencing of purified PCR products and plasmid DNA was performed with BigDye Terminator cycle sequencing kit and a 3730XL genetic analyzer (both from Applied Biosystems, Foster City, CA). The viral sequences were compiled and analyzed with Sequencher software (Gene Codes Corp, Ann Arbor, MI). The following GenBank sequences were used for comparison: for the European (The Netherlands) reference strain (Dumas), X04370 (12); for Oka-P, AB097933 (22); and for Oka-V, AB097932 (22), AF206304 (3), AY016450 (15), and the sequencing information provided by Schmidt et al. (60). Unless otherwise stated, all described nucleotide sequence positions in this paper correspond to the genome of Dumas strain, X04370 (12).

Cloning of PCR products. When direct sequencing did not generate information of sufficient quality or when particular single nucleotide polymorphisms (SNPs) could not be reliably confirmed, additional subcloning was performed, followed by sequencing of numerous generated clones to confirm the consensus sequence of the region. Direct sequencing of the PCR products derived from regions with highly complex secondary structure (flanking regions between internal repeat long and internal repeat short regions, and the R3 repeat region) was complemented by subcloning of amplicons and sequencing of plasmid clones. PCR products containing these regions were individually inserted into a pCR2.1 vector (Invitrogen, Carlsbad, CA) and then transformed into competent *Escherichia coli* by the TOPO TA cloning method (Invitrogen). The plasmid DNAs were purified from cultured bacteria with a QIAprep spin kit (Qiagen). DNA sequences of the cloned inserts were determined using vector-specific sequencing primers.

Sequencing of ends of the viral genomes. The direct sequencing data for viral genome ends were complemented by sequencing of overlapping amplicons generated after circularization using T4 DNA ligase (Roche). The PCR mixtures contained 500 μ M of each deoxynucleoside triphosphate, 10 pmol of each primer, and 2.5 U high-fidelity Platinum Taq polymerase (Invitrogen). PCR products were inserted into a pCR4-TOPO vector and transformed into competent *E. coli* TOP10 bacteria by the TOPO TA cloning method (Invitrogen). The plasmid DNAs were purified with a QIAprep spin kit (Qiagen). The consensus sequence of the cloned amplicons was confirmed by sequencing and alignment of multiple *E. coli* plasmid clones.

Nucleotide sequence accession numbers. The complete nucleotide consensus sequences of the Oka-V_{GSK} (Varilrix) and Oka-V_{Merck} (Varivax) strains are

TABLE 1. Complete Oka-V_{GSK} genome annotation

Start	Stop	Feature ^a	ORF	Function or comment	Start	Stop	Feature ^a	ORF	Function or comment
88	89	Miscellaneous		TRL/UL boundary	65795	65800	Poly(A) signal		
914	587	Gene	1		64743	65768	CDS	36	Thymidine kinase
592	587	Poly(A) signal			65817	65821	3'end of dPyKmRNA		
914	588	CDS	1		66010	68552	Gene	37	
1133	1861	Gene	2		68747	68552	Poly(A) signal		
1133	1849	CDS	2		66010	68535	CDS	37	gH
1856	1861	Poly(A) signal			70229	68583	Gene	38	
2446	1889	Gene	3		68588	68583	Poly(A) signal		
1894	1889	Poly(A) signal			70229	68604	CDS	38	
2446	1907	CDS	3		70569	71305	Gene	39	
4140	2781	Gene	4		71300	71305	Poly(A) signal		
2781	2776	Poly(A) signal			70569	71291	CDS	39	
4140	2782	CDS	4	Transactivator, tegument protein	71476	75699	Gene	40	
5273	4251	Gene	5		75694	75699	Poly(A) signal		
5273	4251	CDS	5	gK	71476	75666	CDS	40	Major nucleocapsid protein
8576	5325	Gene	6		75783	76748	Gene	41	
8606	9398	Gene	7		76743	76748	Poly(A) signal		
9393	9398	Poly(A) signal			75783	76733	CDS	41	
8606	9385	CDS	7		77974	76791	Gene	42	
10666	9425	Gene	8		76786	76791	Poly(A) signal		ORF 45+ORF 42
9430	9425	Poly(A) signal			77974	76787	CDS	42	
10666	9476	CDS	8	Deoxyuridine triphosphatase	78105	80136	Gene	43	
10641	10904	CDS	9A ^b	gN	80131	80136	Poly(A) signal		
11008	11963	Gene	9		78105	80135	CDS	43	
11958	11963	Poly(A) signal			80295	81449	Gene	44	
11008	11916	CDS	9	Syncytium formation, virion protein	81444	81449	Poly(A) signal		
12159	13420	Gene	10		80295	81386	CDS	44	
13415	13420	Poly(A) signal			82529	81474	CDS	45	
12159	13391	CDS	10	Transactivator, tegument protein	82654	83253	CDS	46	
13589	16076	Gene	11		83103	84635	CDS	47	Protein kinase, tegument protein
13936	14196	Repeat region		Reiteration R1	84602	86257	CDS	48	
16071	16076	Poly(A) signal			86161	86429	Gene	49	
13589	16003	CDS	11		86424	86429	Poly(A) signal		
16168	18153	Gene	12		86161	86406	CDS	49	
18695	19350	Gene	13		87807	86466	Gene	50	
19345	19350	Poly(A) signal			86471	86466	Poly(A) signal		
18395	19300	CDS	13		87807	86500	CDS	50	
21067	19296	Gene	14		87806	90313	CDS	51	Origin binding protein
19301	19296	Poly(A) signal			90418	92771	Gene	52	
20526	20851	Repeat region		Reiteration R2	92766	92771	Poly(A) signal		
21067	19385	CDS	14		90418	92733	CDS	52	
22432	21198	Gene	15		93775	92775	Gene	53	
21203	21198	Poly(A) signal			92780	92775	Poly(A) signal		
22432	21212	CDS	15		93775	92780	CDS	53	
23748	22522	Gene	16		95909	93600	CDS	54	
24103	25468	Gene	17		95921	98566	CDS	55	
25463	25468	Poly(A) signal			98493	99280	Gene	56	
24103	25467	CDS	17		99275	99280	Poly(A) signal		
26444	25501	Gene	18		98493	99224	CDS	56	
25506	25501	Poly(A) signal			99548	99309	Gene	57	
26444	25524	CDS	18	Ribonucleotide reductase, small subunit	99314	99309	Poly(A) signal		
28796	26469	Gene	19	Ribonucleotide reductase, big subunit	99548	99333	CDS	57	Cytoplasmic protein
30426	28956	Gene	20		100194	99529	CDS	58	
28961	28956	Poly(A) signal			101141	100224	CDS	59	Uracil-DNA glycosylase
30426	28975	CDS	20		101574	101092	CDS	60	gL, chaperone for gH
30710	33856	Gene	21		104410	102926	Gene	61	
33851	33856	Poly(A) signal			102931	102926	Poly(A) signal		
30710	33826	CDS	21	Nucleocapsid	104410	103007	CDS	61	Transactivator, transrepressor
34034	42341	Gene	22		104849	104850	Miscellaneous		UL/IRL boundary
41405	41470	Repeat region		Reiteration R3	104938	104939	Miscellaneous		IRL/IRS boundary
42336	42341	Poly(A) signal			109061	105065	Gene	62	
34034	42325	CDS	22		105071	105065	Poly(A) signal		
43090	42378	Gene	23		109061	105129	CDS	62	Transactivator, tegument protein
42383	42378	Poly(A) signal			109693	109718	Repeat region		Reiteration 4
43090	42383	CDS	23		110017	110278	Origin of replication		Origin of replication
43973	43163	Gene	24		110507	111359	Gene	63	
43168	43163	Poly(A) signal			111352	111357	Poly(A) signal		
43973	43164	CDS	24		110507	111343	CDS	63	Tegument protein
44570	44083	Gene	25		111491	112072	Gene	64	
44088	44083	Poly(A) signal			112067	112072	Poly(A) signal		
44570	44100	CDS	25		111491	112033	CDS	64	
44458	46125	Gene	26		112571	112107	Gene	65	
46079	47195	Gene	27		112112	112107	Poly(A) signal		
47190	47195	Poly(A) signal			112571	112263	CDS	65	Virion protein
46079	47080	CDS	27		112263	112264	Miscellaneous		IRS/US boundary
50588	46983	Gene	28		112968	114172	Gene	66	
46988	46983	Poly(A) signal			114167	114172	Poly(A) signal		
50588	47004	CDS	28	DNA polymerase	112968	114149	CDS	66	Protein kinase
50809	54460	Gene	29		114427	115523	Gene	67	
54455	54460	Poly(A) signal			115518	115523	Poly(A) signal		
50809	54408	CDS	29	ssDNA binding protein	114427	115491	CDS	67	gI
54587	56899	Gene	30		115739	117652	Gene	68	
56944	59584	Gene	31		117647	117652	Poly(A) signal		
59579	59584	Poly(A) signal			115739	117610	CDS	68	gE
56944	59550	CDS	31	gB, fusogen	117498	117499	Miscellaneous		US/TRS boundary
59703	60150	Gene	32		118266	117690	Gene	69	
60145	60150	Poly(A) signal			117495	117490	Poly(A) signal		
59703	60134	CDS	32	Substrate for ORF 47 kinase	118266	117724	CDS	69	
62074	60245	Gene	33		119250	118400	Gene	70	
60250	60245	Poly(A) signal			118405	118400	Poly(A) signal		
62074	60257	CDS	33	Protease	119250	118414	CDS	70	Tegument protein
63846	62107	Gene	34		119479	119742	Origin of replication		Origin of replication
64689	63913	CDS	35		119921	120066	Repeat region		Reiteration R4
64300	64306	promoter		TATA element	120698	124694	Gene	71	
64321	64325	5'end of dPyKmRNA			124689	124694	Poly(A) signal		
64743	65800	Gene	36		120698	124630	CDS	71	Transactivator, tegument protein

^a CDS, coding sequence; dPyKmRNA, deoxypyrimidine kinase mRNA.

^b ORF was annotated according to the work of Gomi et al. (20)

TABLE 2. Complete Oka-V_{Merck} genome annotation

Start	Stop	Feature ^a	ORF	Function or comment	Start	Stop	Feature ^a	ORF	Function or comment
88	89	Miscellaneous		TRL/UL boundary	65793	65797	Poly(A) signal		
914	587	Gene	1		64741	65765	CDS	36	Thymidine kinase
592	587	Poly(A) signal			65814	65818	3' end of dPyKmRNA		
914	588	CDS	1		66007	68549	Gene	37	
1133	1861	Gene	2		68744	68549	Poly(A) signal		
1133	1849	CDS	2		66007	68532	CDS	37	gH
1856	1861	Poly(A) signal			70226	68580	Gene	38	
2446	1889	Gene	3		68585	68580	Poly(A) signal		
1894	1889	Poly(A) signal			70226	68601	CDS	38	
2446	1907	CDS	3		70566	71302	Gene	39	
4140	2781	Gene	4		71297	71302	Poly(A) signal		
2781	2776	Poly(A) signal			70566	71288	CDS	39	
4140	2782	CDS	4	Transactivator, tegument protein	71473	75696	Gene	40	
5273	4251	Gene	5		75691	75696	Poly(A) signal		
5273	4251	CDS	5	gK	71473	75663	CDS	40	Major nucleocapsid protein
8576	5325	Gene	6		75780	76745	Gene	41	
8606	9398	Gene	7		76740	76745	Poly(A) signal		
9393	9398	Poly(A) signal			75780	76730	CDS	41	
8606	9385	CDS	7		77971	76788	Gene	42	
10666	9425	Gene	8		76783	76788	Poly(A) signal		ORF 45+ORF 42
9430	9425	Poly(A) signal			77971	76784	CDS	42	
10666	9476	CDS	8	Deoxyuridine triphosphatase	78102	80133	Gene	43	
10641	10904	CDS	9A ^b		80128	80133	Poly(A) signal		
11008	11963	Gene	9		78102	80132	CDS	43	
11958	11963	Poly(A) signal			80292	81446	Gene	44	
11008	11916	CDS	9	Syncytium formation, virion protein	81441	81446	Poly(A) signal		
12159	13420	Gene	10		80292	81383	CDS	44	
13415	13420	Poly(A) signal			82526	81471	CDS	45	
12159	13391	CDS	10	Transactivator, tegument protein	82651	83250	CDS	46	
13589	16076	Gene	11		83100	84632	CDS	47	Protein kinase, tegument protein
13936	14196	Repeat region		Reiteration R1	84599	86254	CDS	48	
16071	16076	Poly(A) signal			86158	86426	Gene	49	
13589	16003	CDS	11		86421	86426	Poly(A) signal		
16168	18153	Gene	12		86158	86403	CDS	49	
18695	19350	Gene	13		87804	86463	Gene	50	
19345	19350	Poly(A) signal			86468	86463	Poly(A) signal		
18395	19300	CDS	13		87804	86497	CDS	50	
21067	19296	Gene	14		87803	90310	CDS	51	Origin binding protein
19301	19296	Poly(A) signal			90415	92768	Gene	52	
20526	20851	Repeat region		Reiteration R2	92763	92768	Poly(A) signal		
21067	19385	CDS	14		90415	92730	CDS	52	
22432	21198	Gene	15		93772	92772	Gene	53	
21203	21198	Poly(A) signal			92777	92772	Poly(A) signal		
22432	21212	CDS	15		93772	92777	CDS	53	
23748	22522	Gene	16		95906	93597	CDS	54	
24103	25468	Gene	17		95918	98563	CDS	55	
25463	25468	Poly(A) signal			98490	99277	Gene	56	
24103	25467	CDS	17		99272	99277	Poly(A) signal		
26444	25501	Gene	18		98490	99221	CDS	56	
25506	25501	Poly(A) signal			99545	99306	Gene	57	
26444	25524	CDS	18	Ribonucleotide reductase, small subunit	99311	99306	Poly(A) signal		
28796	26469	Gene	19	Ribonucleotide reductase, big subunit	99545	99330	CDS	57	Cytoplasmic protein
30426	28956	Gene	20		100191	99526	CDS	58	
28961	28956	Poly(A) signal			101138	100221	CDS	59	Uracil-DNA glycosylase
30426	28975	CDS	20		101571	101089	CDS	60	gL, chaperone for gH
30710	33856	Gene	21		104407	102923	Gene	61	
33851	33856	Poly(A) signal			102928	102923	Poly(A) signal		
30710	33826	CDS	21	Nucleocapsid	104407	103004	CDS	61	Transactivator, transrepressor
34034	42341	Gene	22		104846	104847	Miscellaneous		UL/IRL boundary
41405	41470	Repeat region		Reiteration R3	104935	104936	Miscellaneous		IRL/IRS boundary
42336	42341	Poly(A) signal			109058	105062	Gene	62	
34034	42325	CDS	22		105068	105062	Poly(A) signal		
43088	42376	Gene	23		109058	105126	CDS	62	Transactivator, tegument protein
42381	42376	Poly(A) signal			109690	109715	Repeat region		Reiteration 4
43088	42381	CDS	23		110014	110277	Origin of replication		Origin of replication
43971	43161	Gene	24		110506	111356	Gene	63	
43166	43161	Poly(A) signal			111351	111356	Poly(A) signal		
43971	43162	CDS	24		110506	111342	CDS	63	Tegument protein
44568	44081	Gene	25		111490	112067	Gene	64	
44086	44081	Poly(A) signal			112062	112067	Poly(A) signal		
44568	44098	CDS	25		111490	112032	CDS	64	
44456	46123	Gene	26		112566	112102	Gene	65	
46077	47193	Gene	27		112107	112102	Poly(A) signal		
47188	47193	Poly(A) signal			112566	112258	CDS	65	Virion protein
46077	47078	CDS	27		112258	112259	Miscellaneous		IRS/US boundary
50586	46981	Gene	28		112963	114167	Gene	66	
46986	46981	Poly(A) signal			114162	114167	Poly(A) signal		
50586	47002	CDS	28	DNA polymerase	112963	114144	CDS	66	Protein kinase
50807	54458	Gene	29		114422	115518	Gene	67	
54453	54458	Poly(A) signal			115513	115518	Poly(A) signal		
50807	54406	CDS	29	Single-stranded-DNA binding protein	114422	115486	CDS	67	gI
54585	56897	Gene	30		115734	117647	Gene	68	
56942	59582	Gene	31		117642	117647	Poly(A) signal		
59577	59582	Poly(A) signal			115734	117605	CDS	68	gE
56942	59548	CDS	31	gB, fusogen	117490	117491	Miscellaneous		US/TRS boundary
59701	60148	Gene	32		118260	117682	Gene	69	
60143	60148	Poly(A) signal			117687	117682	Poly(A) signal		
59701	60132	CDS	32	Substrate for ORF 47 kinase	118260	117718	CDS	69	
62071	60242	Gene	33		119244	118394	Gene	70	
60247	60242	Poly(A) signal			118399	118394	Poly(A) signal		
62071	60254	CDS	33	Protease	119244	118408	CDS	70	Tegument protein
63843	62104	Gene	34		119473	119736	Origin of replication		Origin of replication
64686	63910	CDS	35		119915	120060	Repeat region		Reiteration R4
64297	64303	Promoter		TATA element	120692	124688	Gene	71	
64321	64325	5' end of dPyKmRNA			124683	124688	Poly(A) signal		
64740	65797	Gene	36		120692	124624	CDS	71	Transactivator, tegument protein

^a CDS, coding sequence; dPyKmRNA, deoxypyrimidine kinase mRNA.

^b ORF was annotated according to the work of Gomi et al. (20).

available in GenBank under the accession numbers DQ008354 and DQ008355, respectively.

RESULTS

Oka-V_{GSK} and Oka-V_{Merck} genome organization. The full-length consensus sequence of Oka-V_{GSK} and Oka-V_{Merck} vaccine strains was essentially determined by bidirectional sequencing of overlapping PCR-amplified fragments. Occasionally, when the amplified region contained SNPs that could not be conclusively resolved, the amplified fragments were subcloned and a consensus sequence was derived from multiple plasmid clones. The obtained sequences were assembled and the complete genomes of the vaccines were annotated using the VZV sequence of the Dumas strain published by Davison and Scott as a template (12). The full annotations for Oka-V_{GSK} and Oka-V_{Merck} are presented in Tables 1 and 2, respectively.

The complete genomes of Oka-V_{GSK} and Oka-V_{Merck} strains are comprised of 124,821 and 124,815 bp, respectively. Like the wild-type Dumas strain and the parental Japanese Oka-V strain, the Oka-V_{GSK} and Oka-V_{Merck} genomes consist of a unique long region flanked by terminal repeat long and internal repeat long inverted repeat regions, as well as a unique short region flanked by internal repeat short (IRS) and terminal repeat short (TRS) inverted repeat regions. An origin of replication was found in both the IRS and TRS regions. Four unique reiteration regions (R1 to R4) were found along the genome, with R4 duplicated in the IRS and TRS regions.

All the open reading frames (ORFs) described for the Dumas VZV strain (12) and the Oka vaccine parental strain (22) were found in the two Oka-derived vaccine strains (Tables 1 and 2). The 72 ORFs predicted to encode proteins were evenly distributed on both DNA strands. Three genes were located within the repeat sequences and were therefore duplicated within the VZV genome, so that ORFs 69 to 71 in the IRS region correspond to ORFs 62 to 64 in the TRS region.

Comparison of Oka strain genomes to the Dumas strain genome. The obtained sequences of Oka-V_{GSK} and Oka-V_{Merck} were aligned with the full-length VZV genomes of Oka-P, Oka-V, and Dumas strains. All sequence differences between the four Oka strains and the Dumas strain are given in Table 3. A total of 326 nucleotide positions displaying differences relative to the genome of Dumas strain (X04370 [12]) were identified. Among these, 228 were common to the four Oka strains, and the remaining 98 were specific to one, two, or three of the Oka strains. Several deletions or insertions were found, but most mutations were substitutions of one nucleotide, i.e., SNPs. Frequently, the original nucleotide was nonetheless preserved, resulting in a mixture of two nucleotides present at the same position (Table 3). Because, to our knowledge, the vaccine strains were never cloned, this is consistent with the existence of multiple viral species that evolved during the attenuation process. Multiple SNPs were found to still contain the original Oka-P-specific nucleotide. This supports the cooperative effect of the overall pattern of nucleotide substitutions in the expression of the attenuation phenotype and, to a lesser extent, the contribution of individual SNPs.

The 98 differences between the Oka-V_{GSK} (124,821 bp), Oka-V_{Merck} (124,815 bp), Oka-P (125,125 bp), and Oka-V (125,078 bp) genomes were found in 25 ORFs (ORFs 1, 2,

6, 9A, 10, 11, 14, 18, 21, 22, 31, 35, 39, 45, 47, 48, 50, 51, 52, 54, 55, 62, 64, and 71), the R1 and R3 repeat regions (in ORFs 11 and 22, respectively), and one origin of replication (Table 3).

The total number of differences between the four Oka strains was determined (Table 4). Of the 98 differences identified, 69 were found between Oka-P and Oka-V, 51 between Oka-V and Oka-V_{GSK}, and 68 between Oka-V and Oka-V_{Merck}. Consequently, Oka-V_{Merck} contains 17 more differences that discriminate it from Oka-V compared with Oka-V_{GSK}.

Although the highest convergence was found for Oka-V_{GSK} and Oka-V_{Merck}, they still had 36 nucleotide differences (Table 4). For 12 of these positions, Oka-V_{Merck} had nucleotides matching the Oka-P strain, whereas the Oka-V_{GSK} strain had only a single position (119683) where the sequence was Oka-P-like. Overall, for the positions in which Oka-V_{GSK} differed from Oka-V_{Merck}, the Oka-V_{GSK} sequence was closer to Oka-V, whereas the Oka-V_{Merck} sequence was closer to Oka-P.

Sixty-nine nucleotide changes between the Oka-V and the Oka-P strains were identified (Table 5). Among these 69 differences, 56 positions in Oka-P were identical to the reference Dumas strain, whereas only 11 positions in Oka-V were identical to the Dumas strain. Identical nucleotides for many of these positions were also present in Oka-V_{GSK} and Oka-V_{Merck}.

To better characterize the observed differences, the substitution spectra were analyzed (Fig. 2). The large majority of mutations were SNPs and only partial, with two different nucleotides at the same position. Compared to Oka-P, transitions (i.e., mutations resulting in substitution of a purine for a purine [A↔G] or a pyrimidine for a pyrimidine [C↔T]) were more frequently (64% to 69%) observed for the Oka-V_{GSK} and Oka-V_{Merck} strains than transversions (i.e., mutations resulting in substitution of a purine for a pyrimidine and vice versa; 13% to 17%). Transversions were more common than insertions or deletions (≤10%). The majority of the identified mutations were silent mutations, either because they were located in intergenic regions or because of the degenerated genetic code. A significant proportion of mutations in intragenic regions (~45%) caused single amino acid substitutions in both the Oka-V_{GSK} and Oka-V_{Merck} strains (Fig. 2). No stop or frameshift mutations were identified. All deletions and insertions either were located in intergenic regions or, when located within coding regions, were multiples of three bases.

Comparison of Oka-V_{GSK} and Oka-V_{Merck} genomes. Sequence differences observed between the Oka-V_{GSK} and Oka-V_{Merck} strains are described in Table 6. Only 36 differences were found throughout the complete genomes (i.e., ~125 kb), three of which were repeated in ORF 62 and its duplicate, ORF 71. These 33 nucleotide unique position changes resulted in 14 amino acid changes, 1 each in ORFs 6, 9A, 10, 31, 39, and 52 and 2 each in ORFs 14, 55, and 62/71 and the R3 repeat region.

Among these 36 position differences between Oka-V_{GSK} and Oka-V_{Merck}, Oka-V_{GSK} had 23 nucleotide sequences identical to Oka-V but only 3 identical to Oka-P. In contrast, Oka-V_{Merck} had 18 positions identical to Oka-P but only 6 identical to Oka-V (Table 6 and Fig. 3).

TABLE 3. Comparison of complete genomic sequences of Dumas and Oka strains of VZV^a

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
A→G	1	X	X	X	X	1	1
G→C	3	X	X	X	X	3	3
Deletion of C (from WT)	109	X	X	X	X	109	109
G→C	178	-	-	X	X	177	177
A→G	236	X	X	X	X	235	235
C→T	262	X	X	X	X	261	261
T→C	560	-	X	X	X	559	559
G→A (ORF 1), N, silent	685	X	X	X	X	684	684
T→T/C (ORF 1), Q, silent	703	-	X	C	C	702	702
T→T/C (ORF 1), P, silent	763	-	X	C	C	762	762
T→C (ORF 1), T→A	789	X	X	X	X	788	788
T→C (ORF 1), Q→R	790	X	X	X	X	789	789
T→C (ORF 1), Q→R	791	X	X	X	X	790	790
C→G (ORF 2), G, silent	1838	-	X	-	-	1837	1837
T→T/C	2515	-	X	C	-	2514	2514
A→G (ORF 4), T, silent	3764	X	X	X	X	3763	3763
C→T (ORF 5), K, silent	4258	X	X	X	X	4257	4257
A→G (ORF 6), S→P	5745	-	X	X	A/G	5744	5744
G→T (ORF 6), H→Q	6853	X	X	X	X	6852	6852
C→A (ORF 6), G→V	7091	X	X	X	X	7090	7090
C→T (ORF 6), P, silent	7753	X	X	X	X	7752	7752
T→C	9460	X	X	X	X	9459	9459
G→A (ORF 8), P→S	10079	X	X	X	X	10078	10078
T→C/T (ORF 9A), W→R	10900	-	X	X	-	10899	10899
T→G (ORF 9), S, silent	11890	X	X	X	X	11889	11889
A→G (ORF 9), T→A	11906	X	X	X	X	11905	11905
C→A (ORF 10), P→H	12188	X	X	X	X	12187	12187
T→C (ORF 10), F→S	12284	X	X	X	X	12283	12283
T→C (ORF 10), F→S	12285	X	X	X	X	12284	12284
C→C/T (ORF 10), A→V	12779	-	X	X	-	12778	12778
T→G (ORF 10), G, silent	13173	X	X	X	X	13172	13172
G→A	13407	X	X	X	X	13406	13406
Deletion (ORF 11, R1), ATTGACGACGAGG GAGAGGCGGAGGA GGGAGAGGCGGA GGAGGGAGAGGC GGAGGAGGGA GAG, IDDEGEAEEG EAEEGEAEEGE	14088	X	X	X	X	14086	14086
Deletion (ORF 11, R1), GCGGAGGAGGAC GCG, AEEDA	14199–213	X	X	X	X	14134–48	14134–48
Insertion (ORF 11, R1), CGCGATCGACGAC GAGGGAGAGGCG GAGGAGGA	14242	X	-	X	X	14164–96	14164–96
T→C	14390	X	X	X	X	14344	14344
C→T (ORF 12), V, silent	17404	X	X	X	X	17358	17358
C→T (ORF 12), L, silent	17834	X	X	X	X	17788	17788
C→T (ORF 12), T, silent	18082	X	X	X	X	18036	18036
G→A (ORF 13), K, silent	18467	X	X	X	X	18421	18421
T→T/C (ORF 14), stop	19431	-	X	-	-	19385	19385
A→G (ORF 14), I, silent	19719	X	X	X	X	19673	19673
T→A (ORF 14), Y→F	20656	X	X	X	X	20610	20610
T→C (ORF 14), T→A	20684	X	X	X	X	20638	20638
C→C/T (ORF 14), K, silent	20703	-	-	X	-	20657	20657
A→T (ORF 14), E→V	20711	X	X	-	-	20665	20665
C→A (ORF 14), C→A	20745	X	X	-	-	20699	20699
T→A (ORF 14), T→S	20753	X	X	X	X	20707	20707

Continued on following page

TABLE 3—Continued

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
C→A (ORF 14), K→N	20787	X	X	C/A	-	20741	20741
C→A (ORF 14), K→N	20829	X	X	C/A	C/A	20783	20783
T→A/T (ORF 14), T→S	20837	-	-	X	X	20791	20791
C→A (ORF 14), K→N	20871	-	-	C/A	C/A	20825	20825
A→T (ORF 14), S→T	20879	-	-	A/T	-	20833	20833
C→A (ORF 14), K→N	20913	-	-	A/C	A/C	20867	20867
T→A (ORF 14), T→S	21005	X	X	X	X	20959	20959
G→A (ORF 15), L, silent	21371	X	X	X	X	21325	21325
G→T (ORF 15), R, silent	21734	X	X	X	X	21688	21688
G→A (ORF 15), S, silent	22311	X	X	X	X	22265	22265
A→G	22504	X	X	X	X	22458	22458
A→G (ORF 16), M→T	22794	X	X	X	X	22748	22748
A→G (ORF 16), F, silent	23294	X	X	X	X	23248	23248
Deletion (ORF 17), dCAT (delS)	24516	X	X	X	X	24469	24469
A→G (ORF 17), T→A	24578	X	X	X	X	24529	24529
C→T (ORF 17), T→M	24654	X	X	X	X	24605	24605
G→A (ORF 17), V→I	25067	X	X	X	X	25018	25018
A→G (ORF 18), N, silent	26125	-	X	X	A/G	26076	26076
A→G (ORF 19), H, silent	27523	X	X	X	X	27474	27474
T→G (ORF 20), G, silent	29201	X	X	X	X	29152	29152
C→T/C (ORF 21), T→I	31732	-	X	-	-	31683	31683
A→G (ORF 21), T→A	32274	X	X	X	X	32225	32225
T→C (ORF 21), H, silent	33722	X	X	X	X	33673	33673
T→C (ORF 21), D, silent	33725	X	X	X	X	33676	33676
T→C (ORF 21), N, silent	33728	X	X	X	X	33679	33679
T→C (ORF 22), V, silent	35543	X	X	X	X	35494	35494
A→G (ORF 22), L, silent	37649	X	X	X	X	37600	37600
A→G (ORF 22), I→V	37902	X	X	X	X	37853	37853
T→C (ORF 22), T, silent	38036	-	-	C/T	C/T	37987	37987
T→C (ORF 22), Y→H	38055	X	X	X	X	38006	38006
A→C (ORF 22), P, silent	38081	X	X	X	X	38032	38032
G→A (ORF 22), E, silent	38177	X	X	X	X	38128	38128
G→T (ORF 22), T, silent	38714	X	X	X	X	38665	38665
C→T (ORF 22), A, silent	38717	X	X	X	X	38668	38668
A→G (ORF 22), R, silent	39023	X	X	X	X	38974	38974
T→T/G (ORF 22), P, silent	39227	-	X	X	-	39178	39178
G→A (ORF 22), Q, silent	39263	X	X	X	X	39214	39214
G→A (ORF 22), R→H	39394	X	X	X	X	39345	39345
A→G (ORF 22), V, silent	39530	X	X	X	X	39481	39481
A→G (ORF 22), Q, silent	40388	X	X	X	X	40339	40339
T→C (ORF 22), P, silent	41057	X	X	X	X	41008	41008
G→A	41452	X	X	X	X	41403	41403
C→T (R3 repeat), A→V	41458	X	X	X	X	41409	41409
G→C (R3 repeat), A→V	41459	X	X	X	X	41410	41410
C→T (R3 repeat), A→V	41476	X	-	X	X	41427	41427
Deletion, GCGCAGCCC	41475–83	-	X	-	-	41426–34	41426–34
G→C (R3 repeat), A→V	41476	X	-	X	X	41427	41427
Deletion, GCGCAGCCC	41484–519	X	X	-	-	41435–70	41435–70
GCGCAGACCGTCC							
AGCCCGCAG							
CCC, AQPAQTVQ							
PAQP							

Continued on following page

Downloaded from <http://jvi.asm.org/> on May 13, 2021 by guest

TABLE 3—Continued

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
C→T (R3 repeat), A→V	41485	-	-	X	-	41436	41436
C→C/T (R3 repeat), A→V	41494	-	-	X	-	41445	41445
A→C (R3 repeat), T→P	41499	-	-	X	X	41450	41450
C→T (ORF 22), T, silent	41618	X	X	X	X	41569	41569
G→A (ORF 22), S→N	41764	X	X	X	X	41715	41715
C→G (ORF 22), Q→E	42069	X	X	X	X	42020	42020
C→T (ORF 22), R, silent	42176	X	X	X	X	42127	42127
A→C (ORF 22), A, silent	42242	X	X	X	X	42193	42193
	42403	Del AAA	Del AA	Ins A	Del A	42355	42353
T→G (ORF 23), S, silent	42476	X	X	X	X	42428	42426
T→C (ORF 24), I→V	43262	X	X	X	X	43214	43212
C→T (ORF 26), C, silent	44835	X	X	X	X	44787	44785
A→G (ORF 28), C→R	47162	X	X	X	X	47114	47112
C→T (ORF 28), L, silent	47940	X	X	X	X	47892	47890
T→C (ORF 28), S→G	48050	X	X	X	X	48002	48000
G→A (ORF 28), T, silent	48825	X	X	X	X	48777	48775
G→A (ORF 28), L, silent	49535	X	X	X	X	49487	49485
C→A (ORF 28), G→C	50081	X	X	X	X	50033	50031
C→T (ORF 29), S, silent	51168	X	X	X	X	51120	51118
A→G (ORF 29), Q, silent	52917	X	X	X	X	52869	52867
A→C (ORF 29), I→L	53482	X	X	X	X	53434	53432
G→A (ORF 29), A→T	53938	X	X	X	X	53890	53888
Deletion (ORF 29), ACA TTTCAGGGTCAA, NISGS	54359–73	X	X	X	X	54310	54308
Deletion, T	54562	X	X	X	X	54498	54496
T→C	54564	X	X	X	X	54500	54498
A→G (ORF 30), P, silent	55820	X	X	X	X	55756	55754
A→C (ORF 31), T→P	57224	X	X	X	X	57160	57158
A→C (ORF 31), A, silent	57301	X	X	X	X	57237	57235
G→T (ORF 31), A, silent	57397	X	X	X	X	57333	57331
A→G (ORF 31), I→V	58595	-	A/G	A/G	X	58531	58529
A→A/G (ORF 31), P, silent	59287	-	X	X	X	59223	59221
Insertion, G	59760	X	X	X	X	59697	59695
Deletion	60278	Del5A	Del5A	Del A	Del AA	60214	60211
A→C	60279	X	X	X	X	60215	60212
C→A (ORF 33), A, silent	60405	X	X	X	X	60341	60338
T→G (ORF 33), Y→S	60781	X	X	X	X	60717	60714
G→A (ORF 33), P→L	61018	X	X	X	X	60954	60951
G→A (ORF 33), P→L	61019	X	X	X	X	60955	60952
T→C (ORF 33), N→G	61201	X	X	X	X	61137	61134
T→C (ORF 33), N→G	61202	X	X	X	X	61138	61135
A→G (ORF 35), A, silent	64067	-	X	X	A/G	64003	64000
A→G (ORF 35), C, silent	64136	X	X	X	X	64072	64069
T→C (ORF 35), P, silent	64259	X	X	X	X	64195	64192
T→C (ORF 35), M→V	64375	X	X	X	X	64311	64308
C→T (ORF 36), A, silent	64989	X	X	X	X	64925	64922
C→T (ORF 36), S→L	65669	X	X	X	X	65605	65602
G→T (ORF 37), L, silent	66646	X	X	X	X	66582	66579
C→T (ORF 37), P→L	66879	X	X	X	X	66815	66812
G→A (ORF 37), R→K	68172	X	X	X	X	68108	68105
A→G (ORF 38), T, silent	69349	X	X	X	X	69285	69282
T→C (ORF 38), S→G	69756	X	X	X	X	69692	69689

Continued on following page

TABLE 3—Continued

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
T→C (ORF 39), M→T	71252	-	X	X	C/T	71188	71185
C→T (ORF 40), V, silent	72997	X	X	X	X	72933	72930
T→C (ORF 40), T, silent	73993	X	X	X	X	73929	73926
C→T (ORF 41), V, silent	76530	X	X	X	X	76466	76463
Deletion, T	78144	X	X	X	X	78079	78076
G→T	80244	X	X	X	X	80179	80176
A→G (ORF 44), N→D	80840	X	X	X	X	80775	80772
C→T (ORF 44), A, silent	81187	X	X	X	X	81122	81119
A→A/G (ORF 45), P, silent	82225	-	X	-	-	82160	82157
G→A/G (ORF 47), E, silent	84091	-	X	-	-	84026	84023
A→G (ORF 47), T, silent	84616	X	X	X	X	84551	84548
G→A (ORF 48), R→H	84983	X	X	X	X	84918	84915
C→T (ORF 48), D, silent	85563	X	X	X	X	85498	85495
A→A/G (ORF 48), T→A	85594	-	-	X	X	85529	85526
C→A (ORF 48), Q→K	86170	X	X	X	X	86105	86102
Deletion, CCTGAT AAAC	86484–93	X	X	X	X	86418	86415
T→G	86556	X	X	X	X	86481	86478
A→A/G (ORF 50), C, silent	87280	-	X	-	-	87205	87202
T→C/T (ORF 50), S→G	87306	-	X	-	-	87231	87228
C→T (ORF 50), S, silent	87841	X	X	X	X	87766	87763
G→T (ORF 51), S, silent	88477	X	X	X	X	88402	88399
A→G (ORF 51), T, silent	89734	-	X	X	-	89659	89656
T→C (ORF 51), T, silent	89905	X	X	X	X	89830	89827
G→T (ORF 51), Q→H	90202	X	X	X	X	90127	90124
T→C (ORF 51), S, silent	90217	X	X	X	X	90142	90139
G→A	90392	X	X	X	X	90317	90314
A→A/G (ORF 52), I→V	90535	-	X	X	-	90460	90457
C→T (ORF 52), G, silent	91191	X	X	X	X	91116	91113
A→G (ORF 52), T→A	92026	X	X	X	X	91951	91948
A→G (ORF 52), T→A	92092	X	X	X	X	92017	92014
A→G (ORF 52), H→R	92375	X	X	X	X	92300	92297
T→C (ORF 53), V, silent	92999	X	X	X	X	92924	92921
T→C (ORF 54), L, silent	94167	-	X	X	T/C	94092	94089
A→G (ORF 54), V, silent	94632	X	X	X	X	94557	94554
A→T (ORF 54), T, silent	94641	X	X	X	X	94566	94563
T→C (ORF 54), G, silent	95241	X	X	X	X	95166	95163
G→A (ORF 54), L, silent	95546	X	X	X	X	95471	95468
T→G (ORF 54), E→D	95601	X	X	X	X	95526	95523
T→C (ORF 55), L, silent	97141	X	X	X	X	97066	97063
T→T/C (ORF 55), V→A	97479	-	-	-	X	97404	97401
C→T (ORF 55), I, silent	97591	X	X	X	X	97516	97513
G→A/G (ORF 55), A→T	97748	-	X	X	X	97673	97670
T→C/T (ORF 55), C→R	97796	-	X	X	-	97721	97718
T→C (ORF 55), G, silent	98437	X	X	X	X	98362	98359
T→C (ORF 56), V, silent	98765	X	X	X	X	98690	98687
A→C (ORF 56), T, silent	98807	X	X	X	X	98732	98729
Deletion (ORF 56), TTC, S	99227–29	X	X	X	X	99148	99145
T→G (ORF 57), H→P	99421	X	X	X	X	99343	99340
A→G (ORF 58), Y, silent	99709	X	X	X	X	99631	99628
C→T (ORF 58), V→I	99981	X	X	X	X	99903	99900
T→A (ORF 58), K→N	100114	X	X	X	X	100036	100033
T→G (ORF 58), N→T	100151	X	X	X	X	100073	100070
A→G	100283	X	X	X	X	100205	100202
A→A/G (ORF 59), L→P	101089	X	X	X	X	101011	101008
C→T (ORF 60), A→T	101331	X	X	X	X	101253	101250

Continued on following page

Downloaded from <http://jvi.asm.org/> on May 13, 2021 by guest

TABLE 3—Continued

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
Insertion (ORF 60), ATC	101623	X	X	X	X	101543–101545	101540–101542
T→C	101886	X	X	X	X	101811	101808
C→T	101991	X	X	X	X	101916	101913
G→A	102192	X	X	X	X	102117	102114
A→G	102203	X	X	X	X	102128	102125
Insertion, TCAAGCTTT	102219	X	X	-	-	102144	102141
AAAAACGTACCCCA							
AACTTAAAACGCTC							
AAATTGCCTTTTGG							
AGGCCTGCCCAACG							
GCCATTATCCCTTG							
GATCTAAGATTGAT							
TTGCGGTAACGTTT							
GCCAA							
C→A	102309	X	X	X	X	102234	102231
A→C	102351	X	X	X	X	102276	102273
A→G	102458	X	X	X	X	102383	102380
T→G	102601	X	X	X	X	102526	102523
T→C	103043	X	X	X	X	102968	102965
A→G	104898	X	X	X	X	104823	104820
C→G	105010	X	X	X	X	104935	104932
T→C	105012	X	X	X	X	104937	104934
T→C	105015	X	X	X	X	104940	104937
T→C	105017	X	X	X	X	104942	104939
Insertion, C	105020	X	-	X	X	104946	104943
Deletion, G	105054	X	X	X	X	104979	104976
Deletion, G	105071	X	X	X	X	104995	104992
Insertion, ACAA	105145	X	X	X	X	105075	105072
A→A/G	105169	-	X	X	X	105097	105094
A→A/G (ORF 62), L→S	105310	-	X	X	X	105238	105235
A→G (ORF 62), G, silent	105312	X	X	X	X	105240	105237
T→C (ORF 62), I→V	105356	-	X	X	T/C	105284	105281
A→G (ORF 62), L→P	105451	X	X	X	X	105379	105376
A→C (ORF 62), S→A	105512	X	X	X	X	105440	105437
A→G (ORF 62), V→A	105544	-	X	X	X	105472	105469
T→C (ORF 62), A, silent	105705	-	X	X	X	105633	105630
T→C (ORF 62), R→G	106262	-	X	X	X	106190	106187
T→C (ORF 62), A, silent	107136	-	X	X	X	107064	107061
C→T (ORF 62), A→T	107165	X	X	X	X	107093	107090
T→C (ORF 62), S→G	107252	-	X	X	X	107180	107177
T→C (ORF 62), R, silent	107307	X	X	X	X	107235	107232
A→A/G (ORF 62), V→A	107599	-	X	X	-	107527	107524
C→A (ORF 62), T, silent	107607	X	X	X	X	107535	107532
T→C (ORF 62), A, silent	107715	X	X	X	X	107643	107640
T→C (ORF 62), P, silent	108111	-	X	X	X	108039	108036
A→G (ORF 62), L, silent	108747	X	X	X	X	108675	108672
A→A/G (ORF 62), M→T	108838	-	X	X	X	108766	108763
G→A (ORF 62), H, silent	108951	X	X	X	X	108879	108876
C→G (ORF 62), A, silent	109044	X	X	X	X	108972	108969
A→A/G	109137	-	X	X	X	109065	109062
A→A/G	109200	-	X	X	-	109128	109125
T→C/T	109546	-	-	X	-	109474	109471
G→T	109654	X	X	X	X	109582	109579
Insertion, CAT	109696	X	X	X	X	109625–109627	109622–109624
Insertion, GGGAGGGG	109907	X	X	-	-	109838	109835
GCGCGGTACCCCGC							
CGATGGGGAGGGG							
GCGCGGTACCCCGC							
CGATGGGGAGGGG							
GCGCGGTACCCCGC							
CGATGGGGAGGGG							
GCGCGGTACCCCGC							
CGATGGGGAGGGG							
GCGCGGTACCCCGC							
CGATGGGGAGGGG							
Insertion, GGGAGGGG	109907	X	-	-	-	109838	109835
GCGCGGTACCCCGC							
CGATGGGGAGGGG							

Continued on following page

TABLE 3—Continued

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
G→A	110003	X	X	X	X	109934	109931
Deletion, G	110058	X	X	X	X	109988	109985
G→A (Ori), -, silent	110112	X	X	X	X	109934	109931
Deletion, AT	110212	-	X	X	-	110142	110140–110141
T→G	110214	-	-	-	X	-	110141
Insertion, ATATAG	110214	X	-	-	-	110142	110141
T→G (Ori)	110216	X	X	X	X	110144	110143
T→G (Ori)	110218	X	X	X	X	110146	110145
T→G (Ori)	110220	X	X	X	X	110148	110147
T→G (Ori)	110222	X	X	X	X	110150	110149
T→G (Ori)	110224	X	X	X	X	110152	110151
T→G (Ori)	110226	X	X	X	X	110154	110153
A→G (Ori)	110232	X	X	X	X	110160	110159
A→G (Ori)	110235	X	X	X	X	110163	110162
Deletion, GC	110378–110379	X	X	X	X	110305	110304
A→G (ORF 63), T, silent	111312	X	X	X	X	110238	110237
A→G (ORF 64), Q→R	111650	-	X	A/G	A/G	111576	111575
T→C (ORF 64), Y→H	112093	X	X	X	X	112019	112018
Deletion/insertion	112128	Del A	Del A	Ins 5a	Ins A	112064–112068	112063
G→A	112198	X	X	X	X	112129	112124
A→G (ORF 66), S, silent	114140	X	X	X	X	114071	114066
G→A (ORF 67), P, silent	115041	X	X	X	X	114072	114967
C→T (ORF 68), T→I	115926	X	X	X	X	115857	115852
C→T	117699	X	X	X	X	117630	117625
Deletion/insertion	117769	Del T	Del T	Ins 5T	Ins T	117701–117705	117696
A→G (ORF 69), Y→H	117804	X	X	X	X	117740	117731
T→C (ORF 69), Q→R	118247	-	X	T/C	T/C	118183	118174
T→C (ORF 70), T, silent	118585	X	X	X	X	118521	118512
Deletion, GC	119518–119519	X	X	X	X	119453	119444
Insertion, CTCTCT	119654	X	X	-	-	119588	119579
T→C (Ori)	119656	X	X	-	-	119590	119581
T→C (Ori)	119665	X	X	X	X	119599	119590
A→C (Ori)	119671	X	X	X	X	119605	119596
A→C (Ori)	119673	X	X	X	X	119607	119598
A→C (Ori)	119675	X	X	X	X	119609	119600
A→C (Ori)	119677	X	-	X	X	119611	119602
Deletion, ATATATAT	119677–119684	-	X	-	-	119611–119618	119602–119609
A→C (Ori)	119679	X	-	X	X	119613	119604
A→C (Ori)	119681	X	-	X	X	119615	119606
A→C (Ori)	119683	X	-	X	A/C	119617	119608
C→T (Ori)	119785	X	X	X	X	119719	119710
Deletion, C	119847	X	X	X	X	119780	119771
C→T,	119894	X	X	X	X	119827	119818
Insertion, TACCGCGCC CCCTCCCCATCGGC GGGGTACCGCGCCC CCTCCCCATCGGCG GGGTACCGCGCCCC CTCCCCATCGGCGG GGTACCGCGCCCC TCCCCATCGGCGGG GTACCGCGCCCCCT CCCCATCGGCGGGG	120135	X	X	-	-	120068	120059
Insertion, TACCGCGCC CCCTCCCCATCGGC GGGG	120135	X	-	-	-	120068	120060
Insertion, GAT	120202	X	X	X	X	120136–120138	120127–120129
C→A	120243	X	X	X	X	120179	120170
A→A/G	120351	-	-	X	-	120287	120278
T→T/C	120697	-	X	X	-	120633	120624
T→T/C	120760	-	X	X	X	120696	120687
G→C (ORF 71), A, silent	120853	X	X	X	X	120789	120780
C→T (ORF 71), H, silent	120946	X	X	X	X	120882	120873
T→C/T (ORF 71), M→T	121059	-	X	X	X	120995	120986
T→C (ORF 71), L, silent	121150	X	X	X	X	121086	121077
A→G (ORF 71), P, silent	121786	-	X	X	X	121722	121713

Continued on following page

Downloaded from <http://jvi.asm.org/> on May 13, 2021 by guest

TABLE 3—Continued

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
A→G (ORF 71), A, silent	122182	X	X	X	X	122118	122109
G→T (ORF 71), T, silent	122290	X	X	X	X	122226	122217
T→C/T (ORF 71), V→A	122298	-	X	X	-	122234	122225
A→G (ORF 71), R, silent	122590	X	X	X	X	122526	122517
A→G (ORF 71), S→G	122645	-	X	X	X	122581	122572
G→A (ORF 71), A→T	122732	X	X	X	X	122668	122659
A→G (ORF 71), A, silent	122761	-	X	X	A/G	122697	122688
A→G (ORF 71), R→G	123635	-	X	X	X	123571	123563
A→G (ORF 71), A, silent	124192	-	X	X	X	124128	124119
T→C (ORF 71), V→A	124353	-	X	X	X	124289	124280
T→G (ORF 71), S→A	124385	X	X	X	X	124321	124312
T→C (ORF 71), L→P	124446	X	X	X	X	124382	124373
A→G (ORF 71), I→V	124541	-	X	X	A/G	124477	124468
T→C (ORF 71), G, silent	124585	X	X	X	X	124521	124512
T→C (ORF 71), L→S	124587	-	T/C	T/C	T/C	124523	124514
T→T/C	124728	-	X	X	X	124664	124655
Insertion, TGTT	124750	X	X	X	X	124687–124690	124678–124681
Deletion, C	124834	X	X	X	X	124773	124764
Deletion, C	124851	X	X	X	X	124789	124780
A→G	124880	NA	NA	X	X	124818	124809
A→G	124882	NA	NA	X	X	124820	124811

^a A partial analysis of 20 of these nucleotide differences was published previously (70). Nucleotide positions within ORFs are indicated, as well as the encoded amino acids. Ori, origin of replication; WT, wild type. X, difference relative to Dumas strain; -, identical nucleotide relative to Dumas strain; NA, not applicable; Del, deletion; Ins, insertion. Where applicable, the resulting codon switch is specified.

DISCUSSION

In this study, we compared the complete genomes of the varicella vaccine strains Oka-V_{GSK} and Oka-V_{Merck}, both derived from the original attenuated Oka-V strain (67). Phylogenetic analyses of these sequences along with 16 other complete VZV genomes were recently reported (50, 69), providing new insight into strain variability (69) and evidence of recombination between major circulating VZV clades (50).

Although VZV is a monotypic virus with a very low rate of interstrain sequence variations (0.061%) compared to other members of the *Herpesviridae* family of viruses (between 0.32% and 3.0% [47]), the sequence analysis of the Oka vaccine strains is not straightforward due to the presence of heterogeneous genomes with distinct sequences (21). Therefore, consensus sequencing provides only an indication of the most prevalent bases for each position. In the present study, we determined the full-length sequences of both Oka-V_{GSK} and Oka-V_{Merck} largely by bidirectional sequencing of overlapping PCR fragments, but when direct sequencing did not generate results of sufficient quality, fragments were subcloned and the

consensus sequence was derived from numerous plasmid clones. All sequences obtained were confirmed on both DNA strands. This approach gave a high-quality assessment of the whole genomes of Oka-V_{GSK} and Oka-V_{Merck}, and this is, to our knowledge, the first published comparative analysis of the complete genomes of these two strains.

Comparison with partial sequencing information published on these strains and the other Oka strains, Oka-P and Oka-V, is shown in Table 7 (3, 22, 32, 59, 60, 63, 69). Argaw et al. sequenced approximately 34 kb from the 3' ends of Oka-V, Oka-P, and Oka-V_{Merck} strains, and Schmidt et al. sequenced approximately 26 kb of the Oka-V_{GSK} strain (3, 60). Two sequence differences were found for Oka-V_{Merck} in ORF 59 (position 101089; A versus A/G) and ORF 62 (position 105310; G versus A/G) (3). Six differences were observed between the present results and those previously published for Oka-V_{GSK}, and 13 differences were observed for Oka-V_{Merck} (60). Finally, comparison between the present study and a previous one (32) revealed quantitative (number of sequence differences between Oka-V_{GSK} and Oka-V_{Merck} strains) and qualitative (ORFs involved) discrepancies.

Our analysis of Oka-V_{GSK} and Oka-V_{Merck} sequences revealed that they have very few nucleotide differences. When we compared their complete genomic sequences (i.e., ~125 kb), we found that only 36 positions were different between Oka-V_{GSK} and Oka-V_{Merck}. These differences lead to 14 unique amino acid substitutions, which suggests that although these two vaccine strains are not identical, they are very similar. The differences resulting in amino acid substitutions were found in 10 different ORFs and in the R3 repeat region, while silent

TABLE 4. Numbers of genomic sequence differences between the four Oka strains

Strain	No. of differences from:		
	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}
Oka-P	69 (Table 5)	79	64
Oka-V		51	68
Oka-V _{GSK}			36 (Table 6)

TABLE 5. Comparison of complete genomic sequences of Oka-P and Oka-V strains of VZV^a

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
T→C	560	-	X	X	X	559	559
T→T/C (ORF 1), Q, silent	703	-	X	C	C	702	702
T→T/C (ORF 1), P, silent	763	-	X	C	C	762	762
C→G (ORF 2), G, silent	1838	-	X	-	-	1837	1837
T→T/C	2515	-	X	C	-	2514	2514
A→G (ORF 6), S→P	5745	-	X	X	A/G	5744	5744
T→C/T (ORF 9A), W→R	10900	-	X	X	-	10899	10899
C→C/T (ORF 10), A→V	12779	-	X	X	-	12778	12778
Insertion (ORF 11, R1), CGCGATCGA CGACGAGGGAGAGCGGAGG AGGA	14242	X	-	X	X	14164–14196	14164–14196
T→T/C (ORF 14), stop	19431	-	X	-	-	19385	19385
A→G (ORF 18), N, silent	26125	-	X	X	A/G	26076	26076
C→T/C (ORF 21), T→I	31732	-	X	-	-	31683	31683
T→T/G (ORF 22), P, silent	39227	-	X	X	-	39178	39178
C→T (R3 repeat), A→V	41476	X	X	X	X	41427	41427
Deletion, GCGCAGCCC	41475–83	-	X	-	-	41426–41434	41426–41434
G→C (R3 repeat), A→V	41476	X	-	X	X	41427	41427
Deletion/insertion	42403	Del AAA	Del AA	Ins A	Del A	42355	42353
A→G (ORF 31), I→V	58595	-	A/G	A/G	X	58531	58529
A→A/G (ORF 31), P, silent	59287	-	X	X	X	59223	59221
A→G (ORF 35), A, silent	64067	-	X	X	A/G	64003	64000
T→C (ORF 39), M→T	71252	-	X	X	C/T	71188	71185
A→A/G (ORF 45), P, silent	82225	-	X	-	-	82160	82157
G→A/G (ORF 47), E, silent	84091	-	X	-	-	84026	84023
A→A/G (ORF 50), C, silent	87280	-	X	-	-	87205	87202
T→C/T (ORF 50), S→G	87306	-	X	-	-	87231	87228
A→G (ORF 51), T, silent	89734	-	X	X	-	89659	89656
A→A/G (ORF 52), I→V	90535	-	X	X	-	90460	90457
T→C (ORF 54), L, silent	94167	-	X	X	T/C	94092	94089
G→A/G (ORF 55), A→T	97748	-	X	X	X	97673	97670
T→C/T (ORF 55), C→R	97796	-	X	X	-	97721	97718
Insertion, C	105020	X	-	X	X	104946	104943
A→A/G	105169	-	X	X	X	105097	105094
A→A/G (ORF 62), L→S	105310	-	X	X	X	105238	105235
T→C (ORF 62), I→V	105356	-	X	X	T/C	105284	105281
A→G (ORF 62), V→A	105544	-	X	X	X	105472	105469
T→C (ORF 62), A, silent	105705	-	X	X	X	105633	105630
T→C (ORF 62), R→G	106262	-	X	X	X	106190	106187
T→C (ORF 62), A, silent	107136	-	X	X	T/C	107064	107061
T→C (ORF 62), S→G	107252	-	X	X	X	107180	107177
A→A/G (ORF 62), V→A	107599	-	X	X	-	107527	107524
T→C (ORF 62), P, silent	108111	-	X	X	X	108039	108036
A→A/G (ORF 62), M→T	108838	-	X	X	X	108766	108763
A→A/G	109137	-	X	X	X	109065	109062
A→A/G	109200	-	X	X	X	109128	109125
Insertion, GGGAGGGGGCGCGGTAC CCCGCCGATG	109907	X	-	-	-	109838	109835
Deletion, AT	110212	-	X	X	-	110142	110140–110141
Insertion, ATATAG	110214	X	-	-	-	110142	110141
A→G (ORF 64), Q→R	111650	-	X	A/G	A/G	111576	111575
T→C (ORF 69), Q→R	118247	-	X	T/C	T/C	118183	118174
A→C (Ori)	119677	X	-	X	X	119611	119602
Deletion, ATATATAT	119677–119684	-	X	-	-	119611–119618	119602–119609
A→C (Ori)	119679	X	-	X	X	119613	119604
A→C (Ori)	119681	X	-	X	X	119615	119606
A→C (Ori)	119683	X	-	X	A/C	119617	119608
Insertion, TACCGCGCCCCCTCCCCA TCGGCGGGG	120135	X	-	-	-	120068	120060
T→T/C	120697	-	X	X	-	120633	120624
T→T/C	120760	-	X	X	X	120696	120687
T→C/T (ORF 71), M→T	121059	-	X	X	X	120995	120986
A→G (ORF 71), P, silent	121786	-	X	X	X	121722	121713
T→C/T (ORF 71), V→A	122298	-	X	X	-	122234	122225
A→G (ORF 71), S→G	122645	-	X	X	X	122581	122572
A→G (ORF 71), A, silent	122761	-	X	X	A/G	122697	122688
A→G (ORF 71), R→G	123635	-	X	X	X	123571	123563
A→G (ORF 71), A, silent	124192	-	X	X	X	124128	124119
T→C (ORF 71), V→A	124353	-	X	X	X	124289	124280
A→G (ORF 71), I→V	124541	-	X	X	A/G	124477	124468
T→C (ORF 71), L→S	124587	-	T/C	T/C	T/C	124523	124514
T→T/C	124728	-	X	X	X	124664	124655

^a Ori, origin of replication; WT, wild type. X, difference relative to Dumas strain; -, identical nucleotide relative to Dumas strain; Del, deletion; Ins, insertion. Where applicable, the resulting codon switch is specified. Boldface highlights homologies between genomic sequences of Oka-V and genomic sequences of Oka-V_{GSK} and/or Oka-V_{Merck}.




FIG. 2. Type (A) and function (B) of the mutations between Oka-P and the Oka-V_{GSK} and Oka-V_{Merck} vaccine strains of VZV. The numbers indicate the number of events identified for each category of mutations. aa, amino acid.

nucleotide substitutions were found in 8 different ORFs, and 1 noncoding substitution was found in the origin of replication.

Transactivation. ORF 62 encodes immediate early protein 62 (IE62), also known as a transcription regulator, which is the major component of the virion tegument and an important transactivating protein for all classes of VZV promoters (28, 49, 56). It is located in the short repeat sequences and has therefore a duplicate gene, ORF 71. These two duplicated genes cover 7% of the whole VZV genome. Recent studies suggested that ORF 62 could play a central role in the attenuated phenotype of the Oka vaccine strains (3, 20–22). Defined amino acid substitutions in ORF 62 that are associated with individual virus variants purified from the vaccine mixture have been linked to enhanced virus growth and spread in monolayer cell culture (22).

The present analysis of vaccine strains confirmed that a high number of mutations could be detected within ORF 62 (20–22). As previously discussed, these SNPs in ORF 62/71 may be important for attenuation of VZV (50, 69). The current analysis identified a nucleotide transition (position 105356 in ORF 62, corresponding to 124541 in ORF 71) that altered an Ile of the IE62 protein to a Val only partially in Oka-V_{Merck} and completely in Oka-V_{GSK} and Oka-V. Because Oka-P encodes only Ile at this position, it is likely that the Oka-V_{Merck} passaging history has selected for minor Oka-P-related species that might be present in the Oka-V vaccine. The second substitution (position 107599 in ORF 62 and 122298 in ORF 71) partially changed a Val to Ala in Oka-V_{GSK} and Oka-V; for Oka-V_{Merck}, this position is identical to the one found in Oka-P and encodes Val only. In both cases, the amino acids involved are small hydrophobic residues. Gomi et al. demonstrated that five amino acid substitutions, including the 105356

mutation, in the carboxyl terminus of IE62 directly reduced transactivational activity (22). Experiments with recombinant VZV will be required to determine how these mutations in ORF 62 modulate VZV gene expression and which amino acid substitutions are responsible for the differences in viral spreading.

The product of ORF 10, the virion-associated transactivator, is a tegument protein that regulates the IE62 promoter (28, 46). A nucleotide substitution (C→C/T) at position 12779 results in a conversion of an Ala in the Dumas, Oka-P, and Oka-V_{Merck} strains to a mixture of Ala and Val in the Oka-V and Oka-V_{GSK} strains. Similarly, minor subspecies from Oka-V that were originally present in Oka-P may have been selected in Oka-V_{Merck}. This position, which corresponds to a location in the middle of the protein, in Oka-V and Oka-V_{GSK} encodes two small hydrophobic amino acids (Val and Ala) that could have similar functions. Indeed, no statistically significant differences in transactivational activity of the ORF 10 gene product could be detected between the wild type and the mutant form, suggesting that this alternative form of ORF 10 has a minimal effect on viral attenuation through modulation of the expression level of IE62 (22). Furthermore, *in vitro* studies have shown that ORF 10 product was dispensable for VZV replication *in vitro* (10).

The helicase-primase complex. The helicase-primase complex consists of three proteins encoded by ORF 6 (primase) and ORFs 52 and 55 (helicase). Interestingly, we found four amino acid substitutions in these proteins, three of which were described previously (22).

The first amino acid substitution was located near the C terminus of ORF 6 (position 5745, A→G), which was a Ser in Dumas and Oka-P, a Pro in Oka-V and Oka-V_{GSK}, and a

TABLE 6. Comparison of complete genomic sequences of Oka-V_{GSK} and Oka-V_{Merck} vaccine strains of VZV^a

Feature relative to WT (Dumas)	Position (WT)	Feature in:				Position in:	
		Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
T→T/C	2515	-	X	C	-	2514	2514
A→G (ORF 6), S→P	5745	-	X	X	A/G	5744	5744
T→C/T (ORF 9A), W→R	10900	-	X	X	-	10899	10899
C→C/T (ORF 10), A→V	12779	-	X	X	-	12778	12778
C→C/T (ORF 14), K, silent	20703	-	-	X	-	20657	20657
C→A (ORF 14), K→N	20787	X	X	C/A	-	20741	20741
A→T (ORF 14), S→T	20879	-	-	A/T	-	20833	20833
A→G (ORF 18), N, silent	26125	-	X	X	A/G	26076	26076
T→T/G (ORF 22), P, silent	39227	-	X	X	-	39178	39178
C→T (R3 repeat), A→V	41485	-	-	X	-	41436	41436
C→C/T (R3 repeat), A→V	41494	-	-	X	-	41445	41445
Deletion/insertion	42403	Del AAA	Del AA	Ins A	Del A	42355	42353
A→G (ORF 31), I→V	58595	-	A/G	A/G	X	58531	58529
Deletion	60278	Del 5A	Del 5A	Del A	Del AA	60214	60211
A→G (ORF 35), A, silent	64067	-	X	X	A/G	64003	64000
T→C (ORF 39), M→T	71252	-	X	X	C/T	71188	71185
A→G (ORF 51), T, silent	89734	-	X	X	-	89659	89656
A→A/G (ORF 52), I→V	90535	-	X	X	-	90460	90457
T→C (ORF 54), L, silent	94167	-	X	X	T/C	94092	94089
T→T/C (ORF 55), V→A	97479	-	-	-	X	97404	97401
T→C/T (ORF 55), C→R	97796	-	X	X	-	97721	97718
T→C (ORF 62), I→V	105356	-	X	X	T/C	105284	105281
T→C (ORF 62), A, silent	107136	-	X	X	T/C	107064	107061
A→A/G (ORF 62), V→A	107599	-	X	X	-	107527	107524
A→A/G	109200	-	X	X	-	109128	109125
T→C/T	109546	-	-	X	-	109474	109471
Deletion, AT	110212	-	X	X	-	110142	110140–110141
T→G	110214	-	-	-	X	-	110141
Deletion/insertion	112128	Del A	Del A	Ins 5a	Ins A	112064–112068	112063
Deletion/insertion	117769	Del T	Del T	Ins 5T	Ins T	117701–117705	117696
A→C (Ori)	119683	X	-	X	A/C	119617	119608
A→A/G	120351	-	-	X	-	120287	120278
T→T/C	120697	-	X	X	-	120633	120624
T→C/T (ORF 71), V→A	122298	-	X	X	-	122234	122225
A→G (ORF 71), A, silent	122761	-	X	X	A/G	122697	122688
A→G (ORF 71), I→V	124541	-	X	X	A/G	124477	124468

^a WT, wild type. X, difference relative to Dumas strain; -, identical nucleotide relative to Dumas strain; Ori, origin of replication; Del, deletion; Ins, insertion. When applicable, the resulting codon switch is specified. Boldface highlights homologies between genomic sequences of Oka-V and genomic sequences of Oka-V_{GSK} and/or Oka-V_{Merck}.

mixture of both in Oka-V_{Merck}. Pro is a rigid residue that could induce substantial changes in the protein conformation. This nucleotide substitution was also comprised in an AluI restriction site. Interestingly, Quinlivan et al. found no differences between Oka-V_{GSK} and Oka-V_{Merck} by AluI restriction analysis: both Oka-V_{GSK} and Oka-V_{Merck} were A/G (\pm AluI), whereas Oka-V was G (+AluI) (52).

The second substitution occurred in ORF 52 (position 90535, A→A/G). In this case, the amino acid residue in Oka-P and Oka-V_{Merck} was Ile, which was partially changed to Val in Oka-V and Oka-V_{GSK}.

The last two substitutions were located in ORF 55. Val at position 97479 was partially replaced by Ala in Oka-V_{Merck}, and Cys at position 97796 was partially replaced by Arg in Oka-V and Oka-V_{GSK}.

Gomi et al. demonstrated that pathogenicity and spreading of VZV were affected by mutations in ORFs 6, 10, and 62, whereas ORFs 52 and 55 did not seem to be important for efficient VZV spreading (22). Because these substitutions were only partial, they could result in the coexistence of different helicase-primase activities resulting from different isomeric

complexes, as shown by restriction fragment length polymorphism analysis (22). For most of these positions Oka-V_{GSK} was similar to Oka-V, whereas Oka-V_{Merck} was similar to Oka-P.

Envelope glycoproteins. VZV produces at least seven glycoproteins, gK, gC, gB, gH, gL, gI, and gE, which are the products of ORFs 5, 14, 31, 37, 60, 67, and 68, respectively (11). Two putative additional glycoproteins were recently described, gN (ORF 9A) and gM (ORF 50) (55, 77). It is known that VZV glycoproteins induce a strong humoral immune response following either natural infection or vaccination with the Oka strain. The SNPs in the nine VZV glycoproteins were reviewed in a recent comparative analysis (64). Some of them are specific to the vaccine strains and thus could be involved in VZV attenuation.

The product of ORF 68, gE, is the most abundant glycoprotein expressed during infection. A single amino acid substitution in this protein was shown to induce the accelerated replication phenotype of the VZV-MSP mutant strain (57). Recently, Grose et al. reported the sequences of two VZV isolates harboring a D150N mutation within ORF 68 (25). That study identified only one mutation, common to all four Oka




FIG. 3. Sequence comparisons of Oka-V_{GSK} and Oka-V_{Merck} with Oka-P and Oka-V strains of VZV. The 36 nucleotide positions that are different in Oka-V_{GSK} and Oka-V_{Merck} vaccines were compared to the sequence of the original vaccine strain Oka-V and its parental virus, Oka-P.

strains (position 115926, C→T), which induced the replacement of a Thr by Ile.

The product of ORF 31, gB, is the second most abundant and immunogenic envelope glycoprotein of VZV after gE. Along with gH and gC, it seems to play a role in the attachment and penetration of viral particles. It was also shown to have important fusogenic properties in the presence of gE and to be associated with cell-to-cell infection (39, 45). An A→G transition at position 58595 induced a conversion of Ile to Val in Oka-V_{Merck} and a mixture of Ile and Val in both Oka-V_{GSK} and Oka-V. Both amino acids are small and hydrophobic, suggesting that this substitution would probably not affect the properties of this glycoprotein.

ORF 14 exhibited one silent replacement and two amino acid substitutions in its product, gC. At position 20787, a co-existence of C and A induced the partial replacement of a Lys by Asn in Oka-V_{GSK}, whereas the original C residue was completely replaced by A in Oka-V_{Merck}, entirely replacing Lys with Asn. Both residues are hydrophilic; however, Lys is basic and Asn is polar, with an uncharged side chain. Because Asn was not detected in either Oka-P or Oka-V, it is likely that this amino acid substitution evolved as a result of additional vaccine passages and might reflect additional cell culture adaptation. The second modification (position 20879, A→T) was found only in the Oka-V_{GSK} strain. It partially changed a Ser to Thr. Both residues are hydrophilic and polar with an uncharged side chain. Grinfeld et al. showed that products of ORF 14 and ORF 67 were dispensable for the establishment of latency in a rat model (24). However, experiments with

SCID-hu mice showed that gC is important for viral tropism in skin cells and that a decrease in gC plays a critical role in attenuation (44). It was also previously shown that expression of gC is dependent on the strain of VZV, with the gC level of Oka-V being much lower than that of wild-type viruses (29, 37).

A nucleotide substitution at position 71252 induced the replacement of a Met by a Thr in the product of ORF 39, one of the two multiply inserted membrane proteins of VZV. This change was complete in Oka-V_{GSK} and Oka-V but only partial in Oka-V_{Merck}. Although both amino acids are neutral, Thr is polar and Met is hydrophobic. This difference may have some effect on the properties of the protein, although these are unclear at present (23).

We also found a mutation in the gN envelope glycoprotein. This glycoprotein is the product of ORF 9A, a newly identified gene positioned closely upstream of the ORF 9 initiation codon (55). Glycoprotein gN is an 87-amino-acid protein whose amino-terminal extremity overlaps with the first nine amino acids of the ORF 8 product, coded on the complementary strand. The observed T-to-T/C shift at position 10900 involves a Trp-to-Arg switch in the very last amino acid of the protein in Oka-V and Oka-V_{GSK}. This change leads to replacement of a hydrophobic amino acid with a large aromatic side chain at the carboxyl terminus of the protein by a hydrophilic basic amino acid, which could alter the membrane topology of the protein or its stability.

R3 repeat region. The R3 reiteration region is located in ORF 22, the longest ORF in VZV. The product of ORF 22 is

TABLE 7. Comparison of Oka-V_{GSK} and Oka-V_{Merck} genomic sequences with previously published Oka genomic sequences⁴⁵

Reference	Position	ORF	Previously reported feature					Feature in present study	
			Dumas	Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
3	84983	48	G	A	A		A	A	
	85563	48	C	T	T		T	T	
	86484		cctgataaac		-		cctgataaac	cctgataaac	
	86556		T		G		G	G	
	87841	50	C		T		T	T	
	88477	51	G		T		T	T	
	89734	51	A		G		G	A	
	89905	51	T		C		C	C	
	90202	51	G		T		T	T	
	90217	51	T		C		C	C	
	90392		G		A		A	A	
	91191	52	C		T		T	T	
	92092	52	A		G		G	G	
	92375	52	A		G		G	G	
	92999	52	T		C		C	C	
	94167	54	T		C		C	T/C	
	94632	54	T		G		G	G	
	94641	54	A		T		T	T	
	95241	54	T	C	C		C	C	
	95546	54	G		A		A	A	
	97141	55	T		C		C	C	
	97470 ^b	55	G		C		G	G	
	97591	55	C		T		T	T	
	97748	55	G		A		A/G	A/G	
	97834 ^c	55	C		T		C	C	
	98437	55	T		C		C	C	
	98765	56	T		C		C	C	
	98807	56	A		C		C	C	
	99227	56	TTC	-	-		TTC	TTC	
	99709	58	A	G	G		G	G	
	99981	58	C	T	T		T	T	
	100114	58	T	A	A		A	A	
	100151	58	T	G	G		G	G	
	100283		A		G		G	G	
	101089	59	A	A	G		A/G	A/G	
	101331	60	C	T	T		T	T	
	101623	60	-	+ATC	+ATC		+ATC	+ATC	
	101886		T		C		C	C	
	101991		C		T		T	T	
	102192		G		A		A	A	
	102203		A		G		G	G	
	102219				+112 bp				
	102309		C		A		A	A	
	102351		A		C		C	C	
	102458		A		G		G	G	
	102601		T		G		G	G	
	103043		T		C		C	C	
	104898		A		G		G	G	
	105010		cctcctct	cctcctct	gcccttacccc		cctcctct	cctcctct	
	105054		G		Del G		Del G	Del G	
	105063 ^d		G		Del G		Del G	Del G	
	105145			+AACA	+AACA		+AACA	+AACA	
	105310	62	A	A	G		A/G	A/G	
	105312	62	A	G	G		G	G	
	105356	62	T	T	C		C	C	
	105451	62	A		G		G	G	
	105512	62	A		C		C	C	
	105544	62	A		G		G	G	
	105705	62	T		C		C	C	
	106262	62	T	T	C		C	C	
	107136	62	T		C		C	C	
	107165	62	C		T		T	T	
	107252	62	T		C		C	C	
	107307	62	T		C		C	C	
	107607	62	C		A		A	A	
	107715	62	T		C		C	C	

Continued on following page

Downloaded from <http://jvi.asm.org/> on May 13, 2021 by guest

TABLE 7—Continued

Reference	Position	ORF	Previously reported feature					Feature in present study	
			Dumas	Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
	108111	62	T		C			C	C
	108747	62	A		G			G	G
	108951	62	G		A			A	A
	109044	62	C		G			G	G
	109694 ^e				+ATC			+CAT	+CAT
	109762				+27 bp			C	C
	110196				Del TA			No del	No del
	110216		(ga) ₉ gg		(ta) ₆ (gc) ₂ aaga		(ta) ₁₆ gag(ga) ₄	Del GC	(ta) ₁₀ (ga) ₉ aaa(ga) ₄
	110378				Del GC			Del GC	Del GC
	111312	63	A		G			G	G
	111650	64	A	A	G		A/G	A/G	A/G
	112093	64	T	C	C		C	C	C
	112130 ^f		-	-	+A ₈		+A ₈	+A ₁₂	+A ₉
	112198		G	A	A		A	A	A
	114140	66	A		G			G	G
	115041	67	G		A			A	A
	115926	68	C		T			T	T
60	1		A	G	G	G	A	G	G
	3		G	C	C	C	G	C	C
	178		G	G	G	C	C	C	C
	560		T	T	C	C	C	C	C
	703 ^g	1	T	T	T	C	C	C	C
	82225 ^h	45	A	A	A	G	A	A	A
	86363	49	A	A	A	T	A	A	A
	87677	50	A	A	A	G	A	A	A
	89734	51	A	A	G	A	A	G	A
	90115 ⁱ	51	A	A	stop	A	A	A	A
	105054		G	-	-	-	G	Del G	Del G
	105071		G	G	-	-	G	Del G	Del G
	105145	Poly(A)	-	AACA	AACA	AACA	-	ACAA	ACAA
	105169		A	A	A/G	A/G	A	A/G	A/G
	105310	62	A	A	A/G	G	A	A/G	A/G
	105356	62	T	T	C	C	T	C	T/C
	105544	62	A	A	G	G	G	G	G
	124353	71	T	T	C	C	C	C	C
	124541	71	A	A	G	G	G	G	A/G
	124587	71	T	T	C/T	C	T	C	C/T
	124728		T	T	C/T	C/T	T	C/T	C/T
	124750	pA-71	-	TGTT	TGTT	TGTT	TGTT	TGTT	TGTT
	124834		C	-	-	-	C	Del C	Del C
	124851		C	-	-	-	C	Del C	Del C
22	560			T	C			C	C
	703	1		T	T/C			C	C
	763	1		T	T/C			C	C
	2515			T	T/C			C	T
	5745	6		A	G			G	A/G
	10900	9A		T	T/C			T/C	T
	12779 ^j	10		T	T/C			T/C	C
	19431	14		T	T/C			T	T
	26125	18		A	G			G	A/G
	31732	21		C	T/C			C	C
	38036 ^k	22		T	T/C			T/C	T/C
	39227	22		T	T/G			T/G	T
	58595	31		A	A/G			A/G	G
	59287	31		A	A/G			A/G	A/G
	64067 ^l	35		A	A/G			G	A/G
	71252 ^m	39		T	T/C			C	T/C
	82225	45		A	A/G			A	A
	84091	47		G	A/G			G	G
	87280	50		A	A/G			A	A
	87306	50		T	T/C			T	T
	89734 ⁿ	51		A	A/G			G	A
	90535	52		A	A/G			A/G	A

Continued on following page

TABLE 7—Continued

Reference	Position	ORF	Previously reported feature				Feature in present study		
			Dumas	Oka-P	Oka-V	Oka-V _{GSK}	Oka-V _{Merck}	Oka-V _{GSK}	Oka-V _{Merck}
	94167	54		T	C			C	T/C
	97748	55		G	A/G			A/G	A/G
	97796	55		T	T/C			T/C	T
	101089 ^p	59		A	A/G			A/G	A/G
	105169			A	A/G			A/G	A/G
	105310	62		A	A/G			A/G	A/G
	105356	62		T	C			C	T/C
	105544	62		A	G			G	G
	105705	62		T	C			C	C
	106262	62		T	C			C	C
	106710 ^p	62		A	A/G			A	A
	107136	62		T	C			C	T/C
	107252	62		T	C			C	C
	107599	62		A	A/G			A/G	A
	107797 ^q	62		A	A/G			A	A
	108111	62		T	C			C	C
	108838	62		A	A/G			A/G	A/G
	109137			A	A/G			A/G	A/G
	109200			A	A/G			A/G	A
	111650 ^r	64		A	A/G			A/G	A/G

^a Boldface highlights differences (3, 60) and homologies (22) between results from published studies and results from the present study. Lowercase indicates insertion; -, missing nucleotide position.

^b Indicated as G in the Oka-V GenBank submission AB097932.

^c Indicated as C in the Oka-V GenBank submission AB097932.

^d In the present alignment, this position is 105071.

^e In the present alignment, this position is 109696.

^f In the present alignment, this position is 112128.

^g Indicated as Y in the Oka-V GenBank submission AB097932.

^h Indicated as R in the Oka-V GenBank submission AB097932.

ⁱ Indicated as A in the Oka-V GenBank submission AB097932.

^j Indicated as C in the Oka-P GenBank submission AB097933.

^k Indicated as T in the Oka-V GenBank submission AB097932.

^l Indicated as G in the Oka-V GenBank submission AB097932.

^m Indicated as C in the Oka-V GenBank submission AB097932.

ⁿ Indicated as G in the Oka-V GenBank submission AB097932.

^o Indicated as R in the Oka-P GenBank submission AB097933.

^p Indicated as A in the Oka-V GenBank submission AB097932.

^q Indicated as A in the Oka-V GenBank submission AB097932.

^r Indicated as A in the Oka-V GenBank submission AB097932.

homologous to the UL36 virion tegument phosphoprotein of herpes simplex virus type 1 (41, 42). R3 is a highly variable region consisting of repeated elements that can vary in number and combination (12, 22), but the impact of this region on the function of ORF 22 phosphoprotein remains mostly unknown because the function of the phosphoprotein itself is poorly understood.

Despite the high variability of the R3 region, no frame-shift mutations were detected in ORF 22, because the repeated elements are present as multiples of 3 bp. Nevertheless, two mutations (positions 41485 and 41494, C→T) in the Oka-V_{GSK} strain were identified, both of which convert Ala into Val, another amino acid with similar properties that probably does not affect the function of the ORF 22 protein.

Attenuation and reactivation. Comparison of the four Oka strain sequences of VZV indicated that Oka-V_{GSK} is genetically closer to Oka-V than Oka-V_{Merck} is but that Oka-V_{Merck} is closer to Oka-P than Oka-V_{GSK} is (reference 50 and the present study). In agreement with these findings, Sauerbrei et al. demonstrated that Oka-V_{Merck} is genetically closer to Oka-P than Oka-V_{GSK} is (59). Nevertheless, it is well established that both vaccine strains are attenuated but remain

strongly immunogenic (43, 73). Recently, Quinlivan et al. suggested an association of particular SNPs in the VZV genome with frequency of vaccine-induced rash (53). Four SNP positions were suggested to contain nucleotides specific to Oka-P in most of the viruses isolated from vaccine rashes. The first one is a silent nucleotide change within ORF 51 (A→G at position 89734). The nucleotide at this position is A in Oka-V_{Merck} and Oka-P but G in Oka-V_{GSK} and Oka-V. The second SNP, at position 105169, contains mixed A/G nucleotides for Oka-V_{Merck}, Oka-V_{GSK}, and Oka-V, whereas the parental Oka-P contains only A. The third SNP, at position 105356, is located within ORF 62. The change from T to C is responsible for an amino acid switch (Ile→Val) for both Oka-V_{GSK} and Oka-V. At the same position, the parental Oka-P contains T (Ile), and Oka-V_{Merck} contains a mixture of T and C (Ile/Val). The last position (nucleotide 107797) was not identified as a SNP in our sequencing data, which agrees with a previous study (22).

Although the genetic basis of Oka-V attenuation has not been determined, Oka-V and Oka-P genomes have nucleotide differences predicted to change amino acids in every class of viral proteins (3, 21). VZV attenuation is a multifactorial phe-

nomenon whose mechanism remains unclear (80), but it is conceivable that mutations of the vaccine genome, in particular, mutations resulting in amino acid modifications, could affect virulence or latency of the vaccine strain. Recently, Peters and coworkers sequenced 11 VZV genomes from different clades, bringing the current number of available full-length VZV sequences to 18 (50, 69). To assess variations that can occur during serial passage in cell culture, these studies included the four Oka strains (Oka-P and the three Oka vaccine strains) and a VZV strain sequenced at passages 5, 22, and 72. As discussed by Tyler et al. (69), the SNPs in ORF 62/71 found in the three Oka vaccine strains and in the VZV strain at high passage level (S628G, R958G, and I1260V in IE62) could be involved in the attenuation of VZV. In addition, it was suggested that other mutations could play a role, particularly those in regions containing ORFs 30 to 55 (69, 78, 80). However, we found that numerous SNPs contain the original Oka-P nucleotide, supporting the idea that the attenuated phenotype is the result of a cooperative effect between several SNPs rather than the result of selected mutations. Therefore, our analysis of SNP importance is aligned with the conclusions of Tyler et al. regarding VZV attenuation (69).

VZV remains latent in sensory-nerve ganglia and can reactivate later, causing herpes zoster. It was suggested that herpes zoster is less common after vaccination, because initial access of Oka-V to neural cells is reduced by limited skin replication or because Oka-V reactivation and secondary viral infection of skin are less efficient, rather than because of an intrinsic attenuation of Oka-V neurotropism (5). Indeed, even though the Oka-V strain of VZV can cause herpes zoster, it seems to reactivate less often than the wild-type VZV even in immunocompromised children (18, 27, 68), and increased incidences of reactivation after vaccination have not been demonstrated in either clinical studies or postmarketing surveillance (33a, 43, 62, 66, 70, 73). In addition, the Oka vaccines have been shown to elicit a strong and protective immune response against varicella (7, 43, 73).

Conclusion. Overall, this study shows that, throughout the entire VZV genome, only 36 nucleotide positions differ between the Oka-V_{GSK} and Oka-V_{Merck} vaccine strains. Analysis of the complete genome of VZV also shows that, genetically, Oka-V_{GSK} is closer to Oka-V and that Oka-V_{Merck} is closer to Oka-P. Although Oka-V_{GSK} and Oka-V_{Merck} exhibit differences, there is a high degree of conservation between these strains at both the nucleotide and amino acid levels. This result supports the clinical data showing that both vaccines are well tolerated and elicit strong immune responses against varicella.

ACKNOWLEDGMENTS

We are grateful to Catherine Arnaudeau-Bégar, Julie Harriague, and Anne Hepburn for their constructive discussions and editorial assistance in the preparation of the manuscript. Technical support from various sequencing staff members is acknowledged.

REFERENCES

1. American Academy of Pediatrics Committee on Infectious Diseases. 1995. Recommendations for the use of live attenuated varicella vaccine. *Pediatrics* **95**:791–796.
2. Arbeter, A. M., S. E. Starr, and S. A. Plotkin. 1986. Varicella vaccine studies in healthy children and adults. *Pediatrics* **78**:748–756.
3. Argaw, T., J. I. Cohen, M. Klutch, K. Lekstrom, T. Yoshikawa, Y. Asano, and P. R. Krause. 2000. Nucleotide sequences that distinguish Oka vaccine from parental Oka and other varicella-zoster virus isolates. *J. Infect. Dis.* **181**:1153–1157.
4. Arvin, A., and A. Gershon. 2006. Control of varicella: why is a two-dose schedule necessary? *Pediatr. Infect. Dis. J.* **25**:475–476.
5. Arvin, A. M. 2001. Varicella vaccine: genesis, efficacy, and attenuation. *Virology* **284**:153–158.
6. Arvin, A. M. 2002. Varilrix (GlaxoSmithKline). *Curr. Opin. Investig. Drugs* **3**:996–999.
7. Asano, Y. 1996. Varicella vaccine: the Japanese experience. *J. Infect. Dis.* **174**(Suppl. 3):S310–S313.
8. Centers for Disease Control and Prevention. 1996. Prevention of varicella: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm. Rep.* **45**:1–36.
9. Chaves, S. S., P. Gargiullo, J. X. Zhang, R. Civen, D. Guris, L. Mascola, and J. F. Seward. 2007. Loss of vaccine-induced immunity to varicella over time. *N. Engl. J. Med.* **356**:1121–1129.
10. Cohen, J. I., and K. Seidel. 1994. Varicella-zoster virus (VZV) open reading frame 10 protein, the homolog of the essential herpes simplex virus protein VP16, is dispensable for VZV replication in vitro. *J. Virol.* **68**:7850–7858.
11. Cole, N. L., and C. Grose. 2003. Membrane fusion mediated by herpesvirus glycoproteins: the paradigm of varicella-zoster virus. *Rev. Med. Virol.* **13**:207–222.
12. Davison, A. J., and J. E. Scott. 1986. The complete DNA sequence of varicella-zoster virus. *J. Gen. Virol.* **67**:1759–1816.
13. de Ory, F., J. M. Echevarría, G. Kafatos, C. Anastassopoulou, N. Andrews, J. Backhouse, G. Berbers, B. Bruckova, D. I. Cohen, H. de Melker, I. Davidkin, G. Gabutti, L. M. Hesketh, K. Johansen, S. Jokinen, L. Jones, A. Linde, E. Miller, J. Mossong, A. Nardone, M. C. Rota, A. Sauerbrei, F. Schneider, Z. Smetana, A. Tischer, A. Tsakris, and R. Vranckx. 2006. European seroepidemiology network 2: standardisation of assays for seroepidemiology of varicella zoster virus. *J. Clin. Virol.* **36**:111–118.
14. D'Hondt, E., E. Berge, G. Colinet, M. Duchene, and J. Peetermans. 1985. Production and quality control of the Oka-strain live varicella vaccine. *Postgrad. Med. J.* **61**:53–56.
15. Faga, B., W. Maury, D. A. Bruckner, and C. Grose. 2001. Identification and mapping of single nucleotide polymorphisms in the varicella-zoster virus genome. *Virology* **280**:1–6.
16. Gail, K., B. Lee, T. Strine, C. Carraher, A. L. Baughman, M. Eaton, J. Montero, and J. Seward. 2002. Outbreak of varicella at a day-care center despite vaccination. *N. Engl. J. Med.* **347**:1909–1915.
17. Gershon, A. A. 2001. The current status of live attenuated varicella vaccine. *Arch. Virol. Suppl.* **2001**:1–6.
18. Gershon, A. A., P. LaRussa, I. Hardy, S. Steinberg, and S. Silverstein. 1992. Varicella vaccine: the American experience. *J. Infect. Dis.* **166**(Suppl. 1):S63–S68.
19. Goh, P., F. S. Lim, H. H. Han, and P. Willems. 2007. Safety and immunogenicity of early vaccination with two doses of tetavalent measles-mumps-rubella-varicella (MMRV) vaccine in healthy children from 9 months of age. *Infection* **35**:326–333.
20. Gomi, Y., T. Imagawa, M. Takahashi, and K. Yamanishi. 2001. Comparison of DNA sequence and transactivation activity of open reading frame 62 of Oka varicella vaccine and its parental viruses. *Arch. Virol. Suppl.* **2001**:49–56.
21. Gomi, Y., T. Imagawa, M. Takahashi, and K. Yamanishi. 2000. Oka varicella vaccine is distinguishable from its parental virus in DNA sequence of open reading frame 62 and its transactivation activity. *J. Med. Virol.* **61**:497–503.
22. Gomi, Y., H. Sunamachi, Y. Mori, K. Nagaiki, M. Takahashi, and K. Yamanishi. 2002. Comparison of the complete DNA sequences of the Oka varicella vaccine and its parental virus. *J. Virol.* **76**:11447–11459.
23. Govero, J., S. Hall, and T. C. Heineman. 2007. Intracellular localization of varicella-zoster virus ORF39 protein and its functional relationship to glycoprotein K. *Virology* **358**:291–302.
24. Grinfeld, E., C. Sadzot-Delvaux, and P. G. Kennedy. 2004. Varicella-zoster virus proteins encoded by open reading frames 14 and 67 are both dispensable for the establishment of latency in a rat model. *Virology* **323**:85–90.
25. Grose, C., S. Tyler, G. Peters, J. Hiebert, G. M. Stephens, W. T. Ruyechan, W. Jackson, J. Storlie, and G. A. Tipples. 2004. Complete DNA sequence analyses of the first two varicella-zoster virus glycoprotein E (D150N) mutant viruses found in North America: evolution of genotypes with an accelerated cell spread phenotype. *J. Virol.* **78**:6799–6807.
26. Hammond, O., Y. Wang, T. Green, J. Antonello, R. Kuhn, C. Motley, P. Stump, B. Rich, N. Chirmule, and R. D. Marchese. 2006. The optimization and validation of the glycoprotein ELISA assay for quantitative varicella-zoster virus (VZV) antibody detection. *J. Med. Virol.* **78**:1679–1687.
27. Hardy, I., A. A. Gershon, S. P. Steinberg, P. LaRussa, et al. 1991. The incidence of zoster after immunization with live attenuated varicella vaccine. A study in children with leukemia. *N. Engl. J. Med.* **325**:1545–1550.
28. Kinchington, P., J. Houghland, A. Arvin, W. Ruyechan, and J. Hay. 1992. The varicella-zoster virus immediate-early protein IE62 is a major component of virus particles. *J. Virol.* **66**:359–366.
29. Kinchington, P. R., P. Ling, M. Pensiero, A. Gershon, J. Hay, and W. T.

- Ruyechan. 1990. A possible role for glycoprotein gpV in the pathogenesis of varicella-zoster virus. *Adv. Exp. Med. Biol.* **278**:83–91.
30. Knuf, M., P. Habermehl, F. Zepp, W. Mannhardt, M. Kuttinig, P. Muttonen, A. Prieler, H. Maurer, H. Bisanz, N. Tornieporth, D. Descamps, and P. Willems. 2006. Immunogenicity and safety of two doses of tetravalent measles-mumps-rubella-varicella vaccine in healthy children. *Pediatr. Infect. Dis. J.* **25**:12–18.
 31. Krahn, D. L., I. Cho, T. Schofield, and R. W. Ellis. 1997. Comparison of gpELISA and neutralizing antibody responses to Oka/Merck live varicella vaccine (Varivax) in children and adults. *Vaccine* **15**:61–64.
 32. Kraiouchkine, N., A. R. Shaw, P. M. Keller, and D. J. Distefano. 2003. Comparison of the complete genome sequences of Varicella Zoster Virus Oka/Merck (Varivax) and Oka/GSK (Varilrix), abstr. G-1652, p. 295. Abstr. 43rd Annu. Intersci. Conf. Antimicrob. Agents Chemother. Chicago, IL, 14 to 17 September 2003.
 33. Krause, P. R., and D. M. Klinman. 1995. Efficacy, immunogenicity, safety, and use of live attenuated chickenpox vaccine. *J. Pediatr.* **127**:518–525.
 - 33a. Kreth, H. W., B. W. Lee, P. Kosuwon, J. Salazar, N. Gloriani-Barzaga, H. L. Bock, F. Meurice. 16 years of global experience with the first refrigerator-stable varicella vaccine (Varilrix™). BioDrugs, in press.
 34. Kuter, B., H. Matthews, H. Shinefield, S. Black, P. Dennehy, B. Watson, K. Reisinger, L. L. Kim, L. Lupinacci, J. Hartzel, and I. Chan. 2004. Ten year follow-up of healthy children who received one or two injections of varicella vaccine. *Pediatr. Infect. Dis. J.* **23**:132–137.
 35. Kuter, B. J., M. L. Brown, J. Hartzel, W. R. Williams, K. A. Eves, S. Black, H. Shinefield, K. S. Reisinger, C. D. Marchant, B. J. Sullivan, M. Thear, S. Klopfer, J. Xu, J. O. Gress, and F. Schodel. 2006. Safety and immunogenicity of a combination measles, mumps, rubella and varicella vaccine (ProQuad). *Hum. Vaccines* **2**:205–214.
 36. Lau, Y. L., S. J. Vessey, I. S. Chan, T. L. Lee, L. M. Huang, C. Y. Lee, T. Y. Lin, B. W. Lee, K. Kwan, S. M. Kasim, C. Y. Chan, K. M. Kaplan, D. J. Distefano, A. L. Harmon, A. Golie, J. Hartzel, J. Xu, S. Li, H. Matthews, J. C. Sadoff, and A. Shaw. 2002. A comparison of safety, tolerability and immunogenicity of Oka/Merck varicella vaccine and VARILRIX in healthy children. *Vaccine* **20**:2942–2949.
 37. Ling, P., P. R. Kinchington, W. T. Ruyechan, and J. Hay. 1991. A detailed analysis of transcripts mapping to varicella zoster virus gene 14 (glycoprotein V). *Virology* **184**:625–635.
 38. Macartney, K. K., P. Beutels, P. McIntyre, and M. A. Burgess. 2005. Varicella vaccination in Australia. *J. Paediatr. Child Health* **41**:544–552.
 39. Maresova, L., T. J. Pasioka, and C. Grose. 2001. Varicella-zoster Virus gB and gE coexpression, but not gB or gE alone, leads to abundant fusion and syncytium formation equivalent to those from gH and gL coexpression. *J. Virol.* **75**:9483–9492.
 40. Marin, M., D. Guris, S. S. Chaves, S. Schmid, and J. F. Seward. 2007. Prevention of varicella: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm. Rep.* **56**:1–40.
 41. McNabb, D. S., and R. J. Courtney. 1992. Analysis of the UL36 open reading frame encoding the large tegument protein (ICP1/2) of herpes simplex virus type 1. *J. Virol.* **66**:7581–7584.
 42. McNabb, D. S., and R. J. Courtney. 1992. Characterization of the large tegument protein (ICP1/2) of herpes simplex virus type 1. *Virology* **190**:221–232.
 43. Meurice, F., J. L. De Bouver, D. Vandevoorde, S. Woods, and H. Bogaeerts. 1996. Immunogenicity and safety of a live attenuated varicella vaccine (Oka/SB Bio) in healthy children. *J. Infect. Dis.* **174**(Suppl. 3):S324–S329.
 44. Moffat, J. F., L. Zerboni, P. R. Kinchington, C. Grose, H. Kaneshima, and A. M. Arvin. 1998. Attenuation of the vaccine Oka strain of varicella-zoster virus and role of glycoprotein C in alpha herpesvirus virulence demonstrated in the SCID-hu mouse. *J. Virol.* **72**:965–974.
 45. Montalvo, E. A., and C. Grose. 1987. Assembly and processing of the disulfide-linked varicella-zoster virus glycoprotein gpII(140). *J. Virol.* **61**:2877–2884.
 46. Moriuchi, H., M. Moriuchi, and J. I. Cohen. 1995. Proteins and cis-acting elements associated with transactivation of the varicella-zoster virus (VZV) immediate-early gene 62 promoter by VZV open reading frame 10 protein. *J. Virol.* **69**:4693–4701.
 47. Muir, W., K. Hawrami, J. Clarke, and J. Breuer. 2001. Investigation of varicella-zoster virus variation by heteroduplex mobility assay, p. 17–25. *In* C. C. Gershon and A. A. Arvin (ed.), *Immunity to and prevention of herpes zoster*. Springer-Verlag, Vienna, Austria.
 48. Nolan, T., P. McIntyre, D. Robertson, and D. Descamps. 2002. Reactogenicity and immunogenicity of a live attenuated tetravalent measles-mumps-rubella-varicella (MMRV) vaccine. *Vaccine* **21**:281–289.
 49. Perera, L. P., J. D. Mosca, W. T. Ruyechan, G. S. Hayward, S. E. Straus, and J. Hay. 1993. A major transactivator of varicella-zoster virus, the immediate-early protein IE62, contains a potent N-terminal activation domain. *J. Virol.* **67**:4474–4483.
 50. Peters, G. A., S. D. Tyler, C. Grose, A. Severini, M. J. Gray, C. Upton, and G. A. Tipples. 2006. A full-genome phylogenetic analysis of varicella-zoster virus reveals a novel origin of replication-based genotyping scheme and evidence of recombination between major circulating clades. *J. Virol.* **80**:9850–9860.
 51. Quian, J., R. Rüttimann, C. Romero, P. Dall’Orso, A. Cerisola, T. Breuer, M. Greenberg, and T. Verstraeten. 2008. Impact of universal varicella vaccination on 1-year-olds in Uruguay: 1997–2005. *Arch. Dis. Child.* **93**:845–850.
 52. Quinlivan, M., A. A. Gershon, S. P. Steinberg, and J. Breuer. 2005. An evaluation of single nucleotide polymorphisms used to differentiate vaccine and wild type strains of varicella-zoster virus. *J. Med. Virol.* **75**:174–180.
 53. Quinlivan, M. L., A. A. Gershon, M. M. Al Bassam, S. P. Steinberg, P. LaRussa, R. A. Nichols, and J. Breuer. 2007. Natural selection for rash-forming genotypes of the varicella-zoster vaccine virus detected within immunized human hosts. *Proc. Natl. Acad. Sci. USA* **104**:208–212.
 54. Robert Koch Institut. 2001. Empfehlungen der Ständigen Impfkommission (STIKO). *Epidemiol. Bull.* **29**:219–223.
 55. Ross, J., M. Williams, and J. I. Cohen. 1997. Disruption of the varicella-zoster virus dUTPase and the adjacent ORF9A gene results in impaired growth and reduced syncytia formation in vitro. *Virology* **234**:186–195.
 56. Sadzot-Delvaux, C., and B. Rentier. 2001. The role of varicella zoster virus immediate-early proteins in latency and their potential use as components of vaccines. *Arch. Virol. Suppl.* **17**:81–89.
 57. Santos, R. A., C. C. Hatfield, N. L. Cole, J. A. Padilla, J. F. Moffat, A. M. Arvin, W. T. Ruyechan, J. Hay, and C. Grose. 2000. Varicella-zoster virus gE escape mutant VZV-MSP exhibits an accelerated cell-to-cell spread phenotype in both infected cell cultures and SCID-hu mice. *Virology* **275**:306–317.
 58. Sauerbrei, A., I. Farber, A. Brandstadt, M. Schacke, and P. Wutzler. 2004. Immunofluorescence test for sensitive detection of varicella-zoster virus-specific IgG: an alternative to fluorescent antibody to membrane antigen test. *J. Virol. Methods* **119**:25–30.
 59. Sauerbrei, A., E. Rubtcova, P. Wutzler, D. S. Schmid, and V. N. Loparev. 2004. Genetic profile of an Oka varicella vaccine virus variant isolated from an infant with zoster. *J. Clin. Microbiol.* **42**:5604–5608.
 60. Schmidt, M., M. Kress, S. Heinemann, and H. Fickenscher. 2003. Varicella-zoster virus isolates, but not the vaccine strain OKA, induce sensitivity to alpha-1 and beta-1 adrenergic stimulation of sensory neurons in culture. *J. Med. Virol.* **70**:S82–S89.
 61. Seward, J. F., B. M. Watson, C. L. Peterson, L. Mascola, J. W. Pelosi, J. X. Zhang, T. J. Maupin, G. S. Goldman, L. J. Tabony, K. G. Brodovicz, A. O. Jumaan, and M. Wharton. 2002. Varicella disease after introduction of varicella vaccine in the United States, 1995–2000. *JAMA* **287**:606–611.
 62. Sharrar, R. G., P. LaRussa, S. A. Galea, S. P. Steinberg, A. R. Sweet, R. M. Keatley, M. E. Wells, W. P. Stephenson, and A. A. Gershon. 2000. The postmarketing safety profile of varicella vaccine. *Vaccine* **19**:916–923.
 63. Shaw, A. R., N. Kraiouchkine, J. Condra, D. Graham, X. Liu, and D. DiStefano. 2001. Genetic differences between OKA varicella vaccines, abstr. 69. 19th Annu. Meet. Eur. Soc. Paediatr. Infect. Dis.
 64. Storie, J., L. Maresova, W. Jackson, and C. Grose. 2008. Comparative analyses of the 9 glycoprotein genes found in wild-type and vaccine strains of varicella-zoster virus. *J. Infect. Dis.* **197**(Suppl. 2):S49–S53.
 65. Straus, S. E., W. Reinhold, H. A. Smith, W. T. Ruyechan, D. K. Henderson, R. M. Blaese, and J. Hay. 1984. Endonuclease analysis of viral DNA from varicella and subsequent zoster infections in the same patient. *N. Engl. J. Med.* **311**:1362–1364.
 66. Takahashi, M. 2001. 25 years’ experience with the Biken Oka strain varicella vaccine: a clinical overview. *Paediatr. Drugs* **3**:285–292.
 67. Takahashi, M., T. Otsuka, Y. Okuno, Y. Asano, and T. Yazaki. 1974. Live vaccine used to prevent the spread of varicella in children in hospital. *Lancet* **ii**:1288–1290.
 68. Takayama, N., M. Takayama, and J. Takita. 2000. Herpes zoster in healthy children immunized with varicella vaccine. *Pediatr. Infect. Dis. J.* **19**:169–170.
 69. Tyler, S. D., G. A. Peters, C. Grose, A. Severini, M. J. Gray, C. Upton, and G. A. Tipples. 2007. Genomic cartography of varicella-zoster virus: a complete genome-based analysis of strain variability with implications for attenuation and phenotypic differences. *Virology* **359**:447–458.
 70. Vassilev, V. 2005. Stable and consistent genetic profile of Oka varicella vaccine virus is not linked with appearance of infrequent breakthrough cases postvaccination. *J. Clin. Microbiol.* **43**:5415–5416.
 71. Vesikari, T., M. Baer, and P. Willems. 2007. Immunogenicity and safety of a second dose of measles-mumps-rubella-varicella vaccine in healthy children aged 5 to 6 years. *Pediatr. Infect. Dis. J.* **26**:153–158.
 72. Vesikari, T., C. Sadzot-Delvaux, B. Rentier, and A. Gershon. 2007. Increasing coverage and efficiency of measles, mumps, and rubella vaccine and introducing universal varicella vaccination in Europe: a role for the combined vaccine. *Pediatr. Infect. Dis. J.* **26**:632–638.
 73. White, C. J., B. J. Kuter, C. S. Hildebrand, K. L. Isganitis, H. Matthews, W. J. Miller, P. J. Provost, R. W. Ellis, R. J. Gerety, and G. B. Calandra. 1991. Varicella vaccine (VARIVAX) in healthy children and adolescents: results from clinical trials, 1987 to 1989. *Pediatrics* **87**:604–610.
 74. Williams, V., A. Gershon, and P. A. Brunell. 1974. Serologic response to varicella-zoster membrane antigens measured by direct immunofluorescence. *J. Infect. Dis.* **130**:669–672.

75. **Wood, M. J.** 2000. History of varicella zoster virus. *Herpes* 7:60–65.
76. **World Health Organization.** 1998. Varicella vaccines. WHO position paper. *Wkly. Epidemiol. Rec.* 73:241–248.
77. **Yamagishi, Y., T. Sadaoka, H. Yoshii, P. Somboonthum, T. Imazawa, K. Nagaike, K. Ozono, K. Yamanishi, and Y. Mori.** 2008. Varicella-zoster virus glycoprotein M homolog is glycosylated, is expressed on the viral envelope, and functions in virus cell-to-cell spread. *J. Virol.* 82:795–804.
78. **Yamanishi, K.** 2008. Molecular analysis of the Oka vaccine strain of varicella-zoster virus. *J. Infect. Dis.* 197(Suppl. 2):S45–S48.
79. **Zepp, F., U. Behre, K. Kindler, K. H. Laakmann, H. Pankow-Culot, W. Mannhardt-Laakmann, F. Beckers, D. Descamps, and P. Willems.** 2007. Immunogenicity and safety of a tetravalent measles-mumps-rubella-varicella vaccine co-administered with a booster dose of a combined diphtheria-tetanus-acellular pertussis-hepatitis B-inactivated poliovirus-Haemophilus influenzae type b conjugate vaccine in healthy children aged 12-23 months. *Eur. J. Pediatr.* 166:857–864.
80. **Zerboni, L., S. Hinchliffe, M. H. Sommer, H. Ito, J. Besser, S. Stamatis, J. Cheng, D. Distefano, N. Kraiouchkine, A. Shaw, and A. M. Arvin.** 2005. Analysis of varicella zoster virus attenuation by evaluation of chimeric parent Oka/vaccine Oka recombinant viruses in skin xenografts in the SCIDhu mouse model. *Virology* 332:337–346.
81. **Zhou, F., R. Harpaz, A. O. Jumaan, C. A. Winston, and A. Shefer.** 2005. Impact of varicella vaccination on health care utilization. *JAMA* 294:797–802.